

Reasoning Under Uncertainty: Variable elimination

Computer Science cpsc322, Lecture 30

(Textbook Chpt 6.4)

~

March, 23, 2009

Lecture Overview

- **Recap Intro Variable Elimination**
- Variable Elimination
 - Simplifications
 - Example
 - Independence
- Where are we?

Bnet Inference: General

- Suppose the variables of the belief network are X_1, \dots, X_n
- Z is the query variable
- $\underline{Y_1 = v_1, \dots, Y_j = v_j}$ are the observed variables (with their values)
- $\underline{Z_1, \dots, Z_k}$ are the remaining variables
- What we want to compute: $\underline{P(Z | Y_1 = v_1, \dots, Y_j = v_j)}$
- We can actually compute: $P(Z, Y_1 = v_1, \dots, Y_j = v_j)$

$$P(Z | Y_1 = v_1, \dots, Y_j = v_j) = \frac{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}{P(Y_1 = v_1, \dots, Y_j = v_j)} = \frac{\overbrace{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}^{\text{Probability of hypothesis}}}{\sum_Z \overbrace{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}^{\text{Probability of evidence}}}$$

Inference with Factors

We can compute $P(Z, Y_1=v_1, \dots, Y_j=v_j)$ by

- expressing the joint as a factor,

$$f(Z, Y_1, \dots, Y_j, Z_1, \dots, Z_k)$$

- assigning $Y_1=v_1, \dots, Y_j=v_j$
- and summing out the variables Z_1, \dots, Z_k

$$P(Z, Y_1 = v_1, \dots, Y_j = v_j) = \sum_{Z_k} \dots \sum_{Z_1} f(Z, Y_1, \dots, Y_j, Z_1, \dots, Z_k) \Big|_{Y_1=v_1, \dots, Y_j=v_j}$$

Variable Elimination Intro (1)

$$P(Z, Y_1 = v_1, \dots, Y_j = v_j) = \sum_{Z_k} \cdots \sum_{Z_1} f(Z, Y_1, \dots, Y_j, Z_1, \dots, Z_k)_{Y_1=v_1, \dots, Y_j=v_j}$$

- Using the chain rule and the definition of a Bnet, we can write $P(X_1, \dots, X_n)$ as

$$\left. \prod_{i=1}^n P(X_i \mid pX_i) \right\}$$

- We can express the joint factor as a product of factors

$$f(Z, Y_1, \dots, Y_j, Z_1, \dots, Z_j) \quad \prod_{i=1}^n f(X_i, pX_i)$$

$$\underbrace{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}_{\sum_{Z_k} \cdots \sum_{Z_1}} = \underbrace{\prod_{i=1}^n f(X_i, pX_i)}_{Y_1=v_1, \dots, Y_j=v_j}$$

Variable Elimination Intro (2)

Inference in belief networks thus reduces to computing “the sums of products....”

$$P(Z, Y_1 = v_1, \dots, Y_j = v_j) = \underbrace{\sum_{Z_k} \dots \sum_{Z_1}}_{\text{1}} \prod_{i=1}^n f(X_i, pX_i) \underbrace{Y_1 = v_1, \dots, Y_j = v_j}_{\text{2}}$$

(4) (3) (1)
 (2)

1. Construct a factor for each conditional probability.
2. In each factor **assign** the observed variables to their observed values.
3. Multiply the factors
4. For each of the other variables $Z_i \in \{Z_1, \dots, Z_k\}$, sum out Z_i

Lecture Overview

- Recap Intro Variable Elimination
- Variable Elimination
 - Simplifications
 - Example
 - Independence
- Where are we?

How to simplify the Computation?

- Assume we have turned the CPTs into factors and performed the assignments

$$\underbrace{f(X_i, pX_i)}_{\text{vars}} \rightarrow f(\underline{\text{varsX}_i})$$

$$f(C, D, G) \underbrace{G = t}_{=} \rightarrow ? f(C, D)$$

$$\sum_{Z_k} \cdots \sum_{Z_1} \prod_{i=1}^n f(X_i, pX_i) \underbrace{Y_1 = v_1, \dots, Y_j = v_j}_{\text{vars}} \rightarrow \sum_{Z_k} \cdots \sum_{Z_1} \prod_{i=1}^n f(\underline{\text{varsX}_i})$$

Let's focus on the basic case, for instance...

$$\sum_A f(C, D) \times f(A, B, D) \times f(A, E) \times f(D)$$

How to simplify: basic case

Let's focus on the basic case.

$$\sum_{Z_1} \prod_{i=1}^n f(\text{varsX}_i)$$

$$\sum_A f(C, D) \times f(A, B, D) \times f(A, E) \times f(D)$$

- How can we compute efficiently?

Factor out those terms that don't involve Z_1 !

$$\left(\prod_{i|Z_1 \notin \text{varsX}_i} f(\text{varsX}_i) \right) \times \left(\sum_{Z_1} \prod_{i|Z_1 \in \text{varsX}_i} f(\text{varsX}_i) \right)$$

do not contain Z_1 *do contain Z_1*

$$\rightarrow f(C, D) \times f(D) \times \sum_A f(A, B, D) \times f(A, E)$$

General case: Summing out variables efficiently

$$\sum_{Z_k} \dots \sum_{Z_1} f_1 \times \dots \times f_h = \sum_{Z_k} \dots \sum_{Z_2} (f_1 \times \dots \times f_i) \left(\sum_{Z_1} f_{i+1} \times \dots \times f_k \right)$$
$$\sum_{Z_k} \dots \sum_{Z_2} f_1 \times \dots \times f_i \times f'$$

Now to sum out a variable Z_2 from a product $f_1 \times \dots \times f_i \times f'$ of factors, again partition the factors into two sets

- F: those that do not contain Z_2
- F: those that do

$$\sum_{Z_k} \dots \sum_{Z_3} \overbrace{\prod^F}^F \sum_{Z_2} \overbrace{\prod^F}^F$$

Analogy with “Computing sums of products”

This simplification is similar to what you can do in basic algebra with *multiplication* and *addition*

- It takes 14 multiplications or additions to evaluate the expression

$$\underline{ab + ac + ad + ae h} + afh +agh.$$

- This expression be evaluated more efficiently....

$$a(b+c+d+h(e+f+g))$$

7 operations

Variable elimination ordering

Is there only one way to simplify? *No*



$$P(G, D=t) = \sum_{A,B,C} f(A, G) f(B, A) f(C, G) f(B, C)$$

$$P(G, D=t) = \underbrace{\sum_A f(A, G)}_{\sum_B f(B, A)} \underbrace{\sum_C f(C, G)}_{f(B, C)}$$

$$P(G, D=t) = \sum_A f(A, G) \sum_C f(C, G) \sum_B f(B, C) f(B, A)$$

Variable elimination algorithm: Summary

$$P(Z, \cancel{Y_1, \dots, Y_j}, \cancel{Z_1, \dots, Z_j})$$

To compute $P(Z | Y_1=v_1, \dots, Y_j=v_j)$:

1. Construct a factor for each conditional probability.
2. Set the observed variables to their observed values.
3. Given an elimination ordering, simplify/decompose sum of products
4. Perform products and sum out Z_i
5. Multiply the remaining factors (all in ? \mathcal{Z})
6. Normalize: divide the resulting factor $f(Z)$ by $\sum_Z f(Z)$.


Lecture Overview

- Recap Intro Variable Elimination
- Variable Elimination
 - Simplifications
 - Example
 - Independence
- Where are we?

Variable elimination example

Compute $P(\underline{G} \mid \underline{H}=h_1)$.

- $\underline{P(G,H)} = \sum_{A,B,C,D,E,F,I} P(A,B,C,D,E,F,G,H,I)$




Variable elimination example

Compute $P(G | H=h_1)$.

- $P(G, H) = \sum_{A,B,C,D,E,F,I} \underline{P(A,B,C,D,E,F,G,H,I)}$

Chain Rule + Conditional Independence:

$\nearrow P(G, H) = \sum_{A,B,C,D,E,F,I} \underline{P(A)} \underline{P(B|A)} \underline{P(C)} \underline{P(D|B,C)} \underline{P(E|C)} \underline{P(F|D)} \underline{P(G|F,E)} \underline{P(H|G)} \underline{P(I|G)}$



Variable elimination example (step1)

Compute $P(G | H=h_1)$.

$$\bullet \quad P(G, H) = \sum_{A,B,C,D,E,F,I} P(A)P(B|A)P(C)P(D|B,C)P(E|C)P(F|D)P(G|F,E)P(H|G)P(I|G)$$

Factorized Representation:

$$P(G, H) = \sum_{A,B,C,D,E,F,I} f_0(A) f_1(B,A) f_2(C) f_3(D,B,C) f_4(E,C) f_5(F, D) f_6(G,F,E) f_7(H,G) f_8(I,G)$$

- $f_0(A)$

- $f_1(B,A)$

- $f_2(C)$

- $f_3(D,B,C)$


- $f_4(E,C)$

- $f_5(F, D)$

- $f_6(G,F,E)$

- $f_7(H,G)$

- $f_8(I,G)$



Variable elimination example (step 2)

Compute $P(G | H=h_1)$.


Previous state:

$$P(G, H) = \sum_{A,B,C,D,E,F,I} f_0(A) f_1(B,A) f_2(C) f_3(D,B,C) f_4(E,C) f_5(F, D) f_6(G,F,E) \underline{f_7(H,G)} f_8(I,G)$$

Observe H :

$$P(G, H=h_1) = \sum_{A,B,C,D,E,F,I} f_0(A) f_1(B,A) f_2(C) f_3(D,B,C) f_4(E,C) f_5(F, D) f_6(G,F,E) \underline{f_9(G)} f_8(I,G)$$

New factor



- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_9(G)$

Variable elimination example (steps 3-4)

Compute $P(G | H=h_1)$.


Previous state:

$$P(G, H) = \sum_{A,B,C,D,E,F,I} f_0(A) f_1(B,A) f_2(C) f_3(D,B,C) f_4(E,C) f_5(F, D) f_6(G,F,E) f_9(G) f_8(I, G)$$

Elimination ordering A, C, E, I, B, D, F:

$$P(G, H=h_1) = \underline{f_9(G)} \sum_F \sum_D f_5(F, D) \sum_B \sum_I f_8(I, G) \sum_E f_6(G, F, E) \sum_C f_2(C) f_3(D, B, C) f_4(E, C) \boxed{\sum_A f_0(A) f_1(B, A)}$$

- $f_0(A)$
- $f_9(G)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$



Variable elimination example(steps 3-4)

Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .


Previous state:

$$P(G, H=h_1) = f_g(G) \sum_F \sum_D f_5(F, D) \sum_B \sum_I f_8(I, G) \sum_E f_6(G, F, E) \sum_C f_2(C) f_3(D, B, C) f_4(E, C) \sum_A f_0(A) f_1(B, A)$$

Eliminate A:

$$P(G, H=h_1) = f_g(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) \sum_I f_8(I, G) \sum_E f_6(G, F, E) \sum_C f_2(C) f_3(D, B, C) f_4(E, C)$$

- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$



- $f_9(G)$
- $f_{10}(B)$

Variable elimination example(steps 3-4)


Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .

Previous state:

$$P(G, H=h_1) = f_g(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) \sum_I f_8(I, G) \sum_E f_6(G, F, E) \sum_C f_2(C) f_3(D, B, C) f_4(E, C)$$

Eliminate C:

$$P(G, H=h_1) = f_g(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) \sum_I f_8(I, G) \sum_E f_6(G, F, E) \boxed{f_{12}(B, D, E)}$$



- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_9(G)$
- $f_{10}(B)$
- $f_{12}(B, D, E)$

Variable elimination example(steps 3-4)

Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .


Previous state:

$$P(G, H=h_1) = f_g(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) \sum_I f_8(I, G) \sum_E f_6(G, F, E) f_{12}(B, D, E)$$

Eliminate E:

$$P(G, H=h_1) = f_g(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) f_{13}(B, D, F, G) \underline{\sum_I f_8(I, G)}$$

- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_9(G)$
- $f_{10}(B)$
- $f_{12}(B, D, E)$
- $f_{13}(B, D, F, G)$



Variable elimination example(steps 3-4)


Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .

Previous state: $P(G, H=h_1) = f_9(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) f_{13}(B, D, F, G) \sum_I f_8(I, G)$

Eliminate I:

$P(G, H=h_1) = f_9(G) f_{14}(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) f_{13}(B, D, F, G)$

- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_9(G)$
- $f_{10}(B)$
- $f_{12}(B, D, E)$
- $f_{13}(B, D, F, G)$
- $f_{14}(G)$




Variable elimination example(steps 3-4)

Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .

Previous state: $P(G, H=h_1) = f_g(G) f_{14}(G) \sum_F \sum_D f_5(F, D) \sum_B f_{10}(B) f_{13}(B, D, F, G)$

Eliminate B:

$P(G, H=h_1) = f_g(G) f_{14}(G) \sum_F \sum_D f_5(F, D) f_{15}(D, F, G)$



- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_9(G)$
- $f_{10}(B)$
- $f_{12}(B, D, E)$
- $f_{13}(B, D, F, G)$
- $f_{14}(G)$
- $f_{15}(D, F, G)$


Variable elimination example(steps 3-4)

Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .

Previous state: $P(G, H=h_1) = f_g(G) f_{14}(G) \sum_F \sum_D f_5(F, D) f_{15}(D, F, G)$

Eliminate D:

$$P(G, H=h_1) = f_g(G) f_{14}(G) \sum_F f_{16}(F, G)$$



- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_9(G)$
- $f_{10}(B)$
- $f_{12}(B, D, E)$
- $f_{13}(B, D, F, G)$
- $f_{14}(G)$
- $f_{15}(D, F, G)$
- $f_{16}(F, G)$

Variable elimination example(steps 3-4)


Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .

Previous state: $P(G, H=h_1) = f_g(G) f_{14}(G) \sum_F f_{16}(F, G)$

Eliminate F:

$$P(G, H=h_1) = f_g(G) f_{14}(G) \underline{f_{17}(G)}$$

- $f_g(G)$
- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_{10}(B)$
- $f_{12}(B, D, E)$
- $f_{13}(B, D, F, G)$
- $f_{14}(G)$
- $f_{15}(D, F, G)$
- $f_{16}(F, G)$
- $f_{17}(G)$




Variable elimination example (step 5)

Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .

Previous state: $P(G, H=h_1) = f_9(G) f_{14}(G) f_{17}(G)$

Multiply remaining factors:

$$P(G, H=h_1) = f_{18}(G)$$



- $f_9(G)$
- $f_{10}(B)$
- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$

Variable elimination example (step 6)


Compute $P(G | H=h_1)$. Elimination ordering A, C, E, I, B, D, F .

Previous state:

$$P(G, H=h_1) = f_{18}(G)$$

Normalize:

$$P(G | H=h_1) = f_{18}(G) / \sum_{g \in \text{dom}(G)} f_{18}(G)$$



- $f_0(A)$
- $f_1(B, A)$
- $f_2(C)$
- $f_3(D, B, C)$
- $f_4(E, C)$
- $f_5(F, D)$
- $f_6(G, F, E)$
- $f_7(H, G)$
- $f_8(I, G)$
- $f_9(G)$
- $f_{10}(B)$
- $f_{12}(B, D, E)$
- $f_{13}(B, D, F, G)$
- $f_{14}(G)$
- $f_{15}(D, F, G)$
- $f_{16}(F, G)$
- $f_{17}(G)$
- $f_{18}(G)$

Lecture Overview

- Recap Intro Variable Elimination
- Variable Elimination
 - Simplifications
 - Example
 - Independence
- Where are we?

Variable elimination and conditional independence

- Variable Elimination looks incredibly painful for large graphs?
- We used conditional independence.....


$$P(X_1 \dots X_n) = \prod P(X_i | \text{parents}(X_i))$$

- Can we use it to make variable elimination simpler?

Yes, all the variables from which the query is conditional independent given the observations can be pruned from the Bnet

VE and conditional independence: Example

- All the variables from which the query is conditional independent given the observations can be pruned from the Bnet



e.g., $P(G | H=v_1, F=v_2, C=v_3)$.



both paths
from G
to D are
blocked

$G \perp\!\!\!\perp$ conditionally
independent from V given
the observed vars
 H, F, C


Learning Goals for today's class

You can:


- Carry out **variable elimination** by using factor representation and using the factor operations.
- Use techniques to simplify variable elimination.



Big Picture: R&R systems



Answering Query under Uncertainty



Next Class

Probability and Time (*TextBook 6.5*)

Course Elements

- Two Practice Exercises on Bnet available. ↗ ↘
- Assignment 4 will be available on Wednesday and due on Apr the 8th (last class).