

# Marginal Independence and Conditional Independence

Computer Science cpsc322, Lecture 26

*(Textbook Chpt 6.1-2)*

Nov, 2013




# Lecture Overview

- Recap with Example
  - Marginalization
  - Conditional Probability
  - Chain Rule ↩



- Bayes' Rule ↩
- Marginal Independence
- Conditional Independence



our most basic and robust form of knowledge about uncertain environments.

# Recap Joint Distribution

$H = \text{True}$     $H = \text{False}$

• 3 binary random variables:  **$P(H, S, F)$**

- **H**  $\text{dom}(H) = \{h, \neg h\}$  has heart disease, does not have...
- **S**  $\text{dom}(S) = \{s, \neg s\}$  smokes, does not smoke
- **F**  $\text{dom}(F) = \{f, \neg f\}$  high fat diet, low fat diet

# Recap Joint Distribution

Joint Prob. Distribution (JPD)

• 3 binary random variables:  $P(H, S, F)$

- **H**  $\text{dom}(H)=\{h, \neg h\}$  has heart disease, does not have...
- **S**  $\text{dom}(S)=\{s, \neg s\}$  smokes, does not smoke
- **F**  $\text{dom}(F)=\{f, \neg f\}$  high fat diet, low fat diet

		f		$\neg f$	
		s	$\neg s$	s	$\neg s$
→ h	.015	.007	.005	.003	
→ $\neg h$	.21	.51	.07	.18	

$2^3 - 1$        $2^k - 1$        $\sum 1$

# Recap Marginalization

$P(H, S, F)$

	<u>f</u>	
	s	$\neg s$
h	.015	.007
$\neg h$	.21	.51

	<u><math>\neg f</math></u>	
	s	$\neg s$
h	.005	.003
$\neg h$	.07	.18

$$P(H, S) = \sum_{x \in \text{dom}(F)} P(H, S, F = x)$$

$P(H, S)?$  →

	<u>s</u>	<u><math>\neg s</math></u>	
h	.02	.01	.03
$\neg h$	.28	.69	.97
$P(S)?$	.3	.7	

$P(H)?$

# Recap Conditional Probability

$P(H,S)$	<b>s</b>	$\neg$ <b>s</b>	$P(H)$
<b>h</b>	.02	.01	.03
$\neg$ <b>h</b>	.28	.69	.97
$P(S)$	.30	.70	

$$P(S | H) = \frac{P(S, H)}{P(H)}$$

$$P(s | \neg h) = \frac{P(s, \neg h)}{P(\neg h)}$$

Two probability distributions for S

$P(S H)$	<b>s</b>	$\neg$ <b>s</b>
$\rightarrow$ <b>h</b>	.666	.333
$\rightarrow$ $\neg$ <b>h</b>	.29	.71

$P(H|S)$   
do this as an exercise

# Recap Conditional Probability (cont.)

$$P(S | H) = \frac{P(S, H)}{P(H)}$$

$$P(S | H, F)$$

$$P(x_1 \dots x_n | Y_1 \dots Y_k)$$

binary

## Two key points we covered in the previous lecture

- We derived this equality from a possible world semantics of probability
- It is not a probability distributions but... *set of prob. distrib.*
- One for each configuration of the conditioning var(s)  
*if conditioned by k binary vars, set  $2^k$  prob. distributions*

# Recap Chain Rule

$$\underline{P(H, S, F)} = P(H) * P(S|H) * P(F|H, S)$$
$$\downarrow$$
$$\cancel{P(H)} * \frac{P(S, H)}{\cancel{P(H)}} * \frac{P(F, H, S)}{\cancel{P(H, S)}}$$

# Bayes Theorem

$$P(S | H) = \frac{P(S, H)}{P(H)}$$

$$P(H | S) = \frac{P(S, H)}{P(S)}$$

Substitute


↓ rewrite

$$P(H | S) P(S) = \underline{P(S, H)}$$

$$P(S | H) = \frac{P(H | S) P(S)}{P(H)}$$



# Lecture Overview

- Recap with Example and Bayes Theorem
  - **Marginal Independence**
  - Conditional Independence
- 

# Do you always need to revise your beliefs?

*No*..... when your knowledge of **Y**'s value doesn't affect your belief in the value of **X**

**DEF.** Random variable **X** is **marginal independent** of random variable **Y** if, for all  $x_i \in \text{dom}(X)$ ,  $y_k \in \text{dom}(Y)$ ,

$$P(X = x_i \mid Y = y_k) = P(X = x_i)$$

# Marginal Independence: Example

- $X$  and  $Y$  are independent iff:  $P(X) = P(X|Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X|Y) = P(X) \quad \text{or} \quad P(Y|X) = P(Y) \quad \text{or} \quad P(X, Y) = P(X) P(Y)$$

- That is new evidence  $Y$  (or  $X$ ) does not affect current belief in  $X$  (or  $Y$ )

- Ex:  $P(\text{Toothache}, \text{Catch}, \text{Cavity}, \text{Weather}) = P(\text{Toothache}, \text{Catch}, \text{Cavity}) P(\text{Weather})$

$|\text{dom}| = 4 \rightarrow$ 

Sunny
Cloudy
Rainy
Snowy

- JPD requiring  $32$  entries is reduced to two smaller ones ( $8$  and  $4$ )

Joint prob. distribution

# In our example are Smoking and Heart Disease marginally Independent ?

What our probabilities are telling us....?


P(H,S)	s	$\neg$ s	P(H)
h	.02	.01	.03
$\neg$ h	.28	.69	.97

$$P(S|H) \stackrel{?}{=} P(S)$$

No!

P(S)	$\rightarrow$	<u>.30</u>	<u>.70</u>
------	---------------	------------	------------

<u>P(S H)</u>	s	$\neg$ s
$\hookrightarrow$ h	.666	.334
$\rightarrow$ $\neg$ h	.29	.71



# Lecture Overview

- Recap with Example
- Marginal Independence
- Conditional Independence

# Conditional Independence


- With marg. Independence, for  $n$  independent random vars,  $O(2^n) \rightarrow O(n)$

$$P(x_1, \dots, x_n) = P(x_1) \times \dots \times P(x_n)$$

- Absolute independence is powerful **but** when you model a particular domain, it is *rare*.....
- Dentistry is a large field with hundreds of variables, few of which are independent (e.g., *Cavity, Heart-disease*).
- What to do?

# Look for weaker form of independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$



- Are *Toothache* and *Catch* marginally independent?

$$P(\downarrow \mid \downarrow) = P(\text{Toothache}) \quad ? \text{ NO}$$

- BUT If have a cavity, does the probability that the probe catches depend on whether I have a toothache? NO

$$(1) P(\text{catch} \mid \text{toothache}, \text{cavity}) = P(\text{catch} \mid \text{cavity})$$

- What if I haven't got a cavity?

$$(2) P(\text{catch} \mid \text{toothache}, \neg \text{cavity}) = P(\text{catch} \mid \neg \text{cavity})$$

- Each is directly caused by the cavity, but neither has a direct effect on the other

# Conditional independence

- In general, *Catch* is conditionally independent of *Toothache* given *Cavity*.

①  $P(\text{Catch} \mid \text{Toothache}, \text{Cavity}) = P(\text{Catch} \mid \text{Cavity})$

- Equivalent statements:

②  $P(\text{Toothache} \mid \text{Catch}, \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity})$

③  $P(\text{Toothache}, \text{Catch} \mid \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity}) P(\text{Catch} \mid \text{Cavity})$

$$P(x, y \mid z) = P(x \mid z) P(y \mid z)$$



# Proof of equivalent statements

①

$$4 \quad P(X|YZ) = P(X|Z) \Rightarrow$$

$$\Rightarrow \textcircled{A} \frac{P(X,Y,Z)}{P(Y,Z)} = \frac{P(X,Z)}{P(Z)} \Rightarrow \textcircled{2}$$

$$\Rightarrow \frac{P(X,Y,Z)}{P(X,Z)} = \frac{P(Y,Z)}{P(Z)} \Rightarrow P(Y|X,Z) = P(Y|Z)$$

$$\begin{aligned} \textcircled{3} \quad P(X,Y|Z) &= \frac{P(X,Y,Z)}{P(Z)} \xRightarrow{\text{from A}} \frac{P(Y,Z) P(X,Z)}{P(Z)} \cdot \frac{1}{P(Z)} \\ &= \frac{P(Y,Z)}{P(Z)} \cdot \frac{P(X,Z)}{P(Z)} = P(Y|Z) \cdot P(X|Z) \end{aligned}$$

# Conditional Independence: Formal Def.

Sometimes, two variables might not be marginally independent. However, they *become* independent after we observe some third variable

**DEF.** Random variable  $X$  is **conditionally independent** of random variable  $Y$  given random variable  $Z$  if, for all  $x_i \in \text{dom}(X)$ ,  $y_k \in \text{dom}(Y)$ ,  $z_m \in \text{dom}(Z)$

$$P(X = x_i \mid Y = y_k, Z = z_m) = P(X = x_i \mid Z = z_m)$$

That is, knowledge of  $Y$ 's value doesn't affect your belief in the value of  $X$ , given a value of  $Z$

# Conditional independence: Use

- Write out full joint distribution using **chain rule**:

$$\begin{aligned} & \mathbf{P}(\mathit{Cavity}, \mathit{Catch}, \mathit{Toothache}) \\ &= \mathbf{P}(\mathit{Toothache} \mid \mathit{Catch}, \mathit{Cavity}) \mathbf{P}(\mathit{Catch} \mid \mathit{Cavity}) \mathbf{P}(\mathit{Cavity}) \\ &= \mathbf{P}(\mathit{Toothache} \mid \mathit{Cavity}) \mathbf{P}(\mathit{Catch} \mid \mathit{Cavity}) \mathbf{P}(\mathit{Cavity}) \end{aligned}$$

Handwritten annotations: A blue box surrounds the first equation. A blue arrow points from the box to the second equation. A blue bracket under the first term of the second equation is labeled '2'. A blue bracket under the second term is labeled '2'. A blue bracket under the third term is labeled '1'. A blue arrow points from the '2' under the first term to the calculation  $2^3 - 1 = 7$ . Another blue arrow points from the '2' under the second term to the same calculation. A blue arrow points from the '1' under the third term to the calculation  $2 + 2 + 1 = 5$ .

how many probabilities?

$$2^3 - 1 = 7$$

$$2 + 2 + 1 = 5$$


- The use of conditional independence often reduces the size of the representation of the joint distribution from **exponential in  $n$**  to **linear in  $n$** . **What is  $n$ ?** # of vars
- Conditional independence** is our **most basic** and **robust** form of **knowledge** about **uncertain environments**.

# Conditional Independence Example 2

- Given whether there is/isn't power in wire  $w_0$ , is whether light  $l_1$  is lit or not, independent of the position of switch  $s_2$ ?

$$P(l_1 | s_2, w_0) \stackrel{?}{=} P(l_1 | w_0)$$

yes!




# Conditional Independence Example 3

- Is every other variable in the system independent of whether light  $l_1$  is lit, given whether there is power in wire  $w_0$ ?

$$P(s_1 | l_1, w_0) = P(s_1 | w_0)$$

$w_1$   
 $w_2$   
⋮



yes!



# Learning Goals for today's class

- **You can:**
- **Derive the Bayes Rule**
- **Define and use Marginal Independence**
- **Define and use Conditional Independence**

# Where are we? (Summary)

- Probability is a rigorous formalism for uncertain knowledge
- **Joint probability distribution**  specifies probability of every **possible world**
- Queries can be answered by summing over possible worlds
- For nontrivial domains, we must find a way to reduce the joint distribution size
- **Independence** (*rare*) and **conditional independence**  (*frequent*) provide the tools

# Next Class

- Bayesian Networks (Chpt 6.3)

**Start working on assignments3 !**