

Bayesian Games

CPSC 532L Week 6

Lecture Overview

- 1 Quiz
- 2 Bayesian Game Forms
- 3 Analyzing Bayesian games

Lecture Overview

- 1 Quiz
- 2 Bayesian Game Forms**
- 3 Analyzing Bayesian games

Fun Game

- Choose a phone number none of your neighbours knows; consider it to be ABC-DEFG

Fun Game

- Choose a phone number none of your neighbours knows; consider it to be ABC-DEFG
 - take “DE” as your valuation
 - play a first-price auction with three neighbours, where your utility is your valuation minus the amount you pay

Fun Game

- Choose a phone number none of your neighbours knows; consider it to be ABC-DEFG
 - take “DE” as your valuation
 - play a first-price auction with three neighbours, where your utility is your valuation minus the amount you pay
 - now play the auction again, same neighbours, same valuation

Fun Game

- Choose a phone number none of your neighbours knows; consider it to be ABC-DEFG
 - take “DE” as your valuation
 - play a first-price auction with three neighbours, where your utility is your valuation minus the amount you pay
 - now play the auction again, same neighbours, same valuation
 - now play again, with “FG” as your valuation

Fun Game

- Choose a phone number none of your neighbours knows; consider it to be ABC-DEFG
 - take “DE” as your valuation
 - play a first-price auction with three neighbours, where your utility is your valuation minus the amount you pay
 - now play the auction again, same neighbours, same valuation
 - now play again, with “FG” as your valuation

Fun Game

- Choose a phone number none of your neighbours knows; consider it to be ABC-DEFG
 - take “DE” as your valuation
 - play a first-price auction with three neighbours, where your utility is your valuation minus the amount you pay
 - now play the auction again, same neighbours, same valuation
 - now play again, with “FG” as your valuation
- Questions:
 - what is the role of uncertainty here?
 - can we model this uncertainty using an imperfect information extensive form game?

Fun Game

- Choose a phone number none of your neighbours knows; consider it to be ABC-DEFG
 - take “DE” as your valuation
 - play a first-price auction with three neighbours, where your utility is your valuation minus the amount you pay
 - now play the auction again, same neighbours, same valuation
 - now play again, with “FG” as your valuation
- Questions:
 - what is the role of uncertainty here?
 - can we model this uncertainty using an imperfect information extensive form game?
 - imperfect info means not knowing what node you're in in the info set

Definition 1: Information Sets

- **Bayesian game**: a set of games that differ only in their payoffs, a common prior defined over them, and a partition structure over the games for each agent.

Definition (Bayesian Game: Information Sets)

A **Bayesian game** is a tuple (N, G, P, I) where

- N is a set of agents,
- G is a set of games with N agents each such that if $g, g' \in G$ then for each agent $i \in N$ the strategy space in g is identical to the strategy space in g' ,
- $P \in \Pi(G)$ is a common prior over games, where $\Pi(G)$ is the set of all probability distributions over G , and
- $I = (I_1, \dots, I_N)$ is a set of partitions of G , one for each agent.

Definition 1: Example

	$I_{2,1}$	$I_{2,2}$																
$I_{1,1}$	<table style="border-collapse: collapse; margin: auto;"> <tr> <td colspan="2" style="padding: 5px;">MP</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">2, 0</td> <td style="border: 1px solid black; padding: 5px;">0, 2</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">0, 2</td> <td style="border: 1px solid black; padding: 5px;">2, 0</td> </tr> <tr> <td colspan="2" style="padding: 5px;">$p = 0.3$</td> </tr> </table>	MP		2, 0	0, 2	0, 2	2, 0	$p = 0.3$		<table style="border-collapse: collapse; margin: auto;"> <tr> <td colspan="2" style="padding: 5px;">PD</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">2, 2</td> <td style="border: 1px solid black; padding: 5px;">0, 3</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">3, 0</td> <td style="border: 1px solid black; padding: 5px;">1, 1</td> </tr> <tr> <td colspan="2" style="padding: 5px;">$p = 0.1$</td> </tr> </table>	PD		2, 2	0, 3	3, 0	1, 1	$p = 0.1$	
MP																		
2, 0	0, 2																	
0, 2	2, 0																	
$p = 0.3$																		
PD																		
2, 2	0, 3																	
3, 0	1, 1																	
$p = 0.1$																		
$I_{1,2}$	<table style="border-collapse: collapse; margin: auto;"> <tr> <td colspan="2" style="padding: 5px;">Coord</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">2, 2</td> <td style="border: 1px solid black; padding: 5px;">0, 0</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">0, 0</td> <td style="border: 1px solid black; padding: 5px;">1, 1</td> </tr> <tr> <td colspan="2" style="padding: 5px;">$p = 0.2$</td> </tr> </table>	Coord		2, 2	0, 0	0, 0	1, 1	$p = 0.2$		<table style="border-collapse: collapse; margin: auto;"> <tr> <td colspan="2" style="padding: 5px;">BoS</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">2, 1</td> <td style="border: 1px solid black; padding: 5px;">0, 0</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">0, 0</td> <td style="border: 1px solid black; padding: 5px;">1, 2</td> </tr> <tr> <td colspan="2" style="padding: 5px;">$p = 0.4$</td> </tr> </table>	BoS		2, 1	0, 0	0, 0	1, 2	$p = 0.4$	
Coord																		
2, 2	0, 0																	
0, 0	1, 1																	
$p = 0.2$																		
BoS																		
2, 1	0, 0																	
0, 0	1, 2																	
$p = 0.4$																		

Definition 2: Epistemic Types

- Directly represent uncertainty over utility function using the notion of **epistemic type**.

Definition

A **Bayesian game** is a tuple (N, A, Θ, p, u) where

- N is a set of agents,
- $A = (A_1, \dots, A_n)$, where A_i is the set of actions available to player i ,
- $\Theta = (\Theta_1, \dots, \Theta_n)$, where Θ_i is the type space of player i ,
- $p : \Theta \rightarrow [0, 1]$ is the common prior over types,
- $u = (u_1, \dots, u_n)$, where $u_i : A \times \Theta \rightarrow \mathbb{R}$ is the utility function for player i .

Definition 2: Example

	$I_{2,1}$	$I_{2,2}$								
$I_{1,1}$	<p>MP</p> <table border="1"> <tr><td>2, 0</td><td>0, 2</td></tr> <tr><td>0, 2</td><td>2, 0</td></tr> </table> <p>$p = 0.3$</p>	2, 0	0, 2	0, 2	2, 0	<p>PD</p> <table border="1"> <tr><td>2, 2</td><td>0, 3</td></tr> <tr><td>3, 0</td><td>1, 1</td></tr> </table> <p>$p = 0.1$</p>	2, 2	0, 3	3, 0	1, 1
2, 0	0, 2									
0, 2	2, 0									
2, 2	0, 3									
3, 0	1, 1									
$I_{1,2}$	<p>Coord</p> <table border="1"> <tr><td>2, 2</td><td>0, 0</td></tr> <tr><td>0, 0</td><td>1, 1</td></tr> </table> <p>$p = 0.2$</p>	2, 2	0, 0	0, 0	1, 1	<p>BoS</p> <table border="1"> <tr><td>2, 1</td><td>0, 0</td></tr> <tr><td>0, 0</td><td>1, 2</td></tr> </table> <p>$p = 0.4$</p>	2, 1	0, 0	0, 0	1, 2
2, 2	0, 0									
0, 0	1, 1									
2, 1	0, 0									
0, 0	1, 2									

a_1	a_2	θ_1	θ_2	u_1	u_2
U	L	$\theta_{1,1}$	$\theta_{2,1}$	2	0
U	L	$\theta_{1,1}$	$\theta_{2,2}$	2	2
U	L	$\theta_{1,2}$	$\theta_{2,1}$	2	2
U	L	$\theta_{1,2}$	$\theta_{2,2}$	2	1
U	R	$\theta_{1,1}$	$\theta_{2,1}$	0	2
U	R	$\theta_{1,1}$	$\theta_{2,2}$	0	3
U	R	$\theta_{1,2}$	$\theta_{2,1}$	0	0
U	R	$\theta_{1,2}$	$\theta_{2,2}$	0	0

a_1	a_2	θ_1	θ_2	u_1	u_2
D	L	$\theta_{1,1}$	$\theta_{2,1}$	0	2
D	L	$\theta_{1,1}$	$\theta_{2,2}$	3	0
D	L	$\theta_{1,2}$	$\theta_{2,1}$	0	0
D	L	$\theta_{1,2}$	$\theta_{2,2}$	0	0
D	R	$\theta_{1,1}$	$\theta_{2,1}$	2	0
D	R	$\theta_{1,1}$	$\theta_{2,2}$	1	1
D	R	$\theta_{1,2}$	$\theta_{2,1}$	1	1
D	R	$\theta_{1,2}$	$\theta_{2,2}$	1	2

Fun Game 2: Chicken... after dark!

	G	S
G	$\theta_1 - \theta_2 - 2, \theta_2 - \theta_1 - 2$	$3, -2$
S	$-2, 3$	$-1, -1$

- Write down the numbers 0,1,2,10 on a individual pieces of paper. This is the deck of cards.
- Each player draws 1 card (the size/power of your car).
- Play chicken! If you collide, each player's utility depends on the size of both cars.

Fun Game 2: Chicken... after dark!


	G	S
G	$\theta_1 - \theta_2 - 2, \theta_2 - \theta_1 - 2$	$3, -2$
S	$-2, 3$	$-1, -1$

- Write down the numbers 0,1,2,10 on a individual pieces of paper. This is the deck of cards.
- Each player draws 1 card (the size/power of your car).
- Play chicken! If you collide, each player's utility depends on the size of both cars.
- This game is a bit like poker. What's missing?

Definition 3: Extensive Form with Chance Moves

- Add an agent, “Nature,” who follows a commonly known mixed strategy.
- Thus, reduce Bayesian games to extensive form games of imperfect information.
- This definition is cumbersome for the same reason that IIEF is a cumbersome way of representing matrix games like Prisoner’s dilemma
 - however, it makes sense when the agents really do move sequentially, and at least occasionally observe each other’s actions.

Definition 3: Example



Lecture Overview

- 1 Quiz
- 2 Bayesian Game Forms
- 3 Analyzing Bayesian games**

Strategies

- **Pure strategy:** $s_i : \Theta_i \rightarrow A_i$
 - a mapping from every type agent i could have to the action he would play if he had that type.
- **Mixed strategy:** $s_i : \Theta_i \rightarrow \Pi(A_i)$
 - a mapping from i 's type to a probability distribution over his action choices.
- $s_j(a_j|\theta_j)$
 - the probability under mixed strategy s_j that agent j plays action a_j , given that j 's type is θ_j .

Expected Utility

Three meaningful notions of expected utility:

- *ex-ante*
 - the agent knows nothing about anyone's actual type;
- *ex-interim*
 - an agent knows his own type but not the types of the other agents;
- *ex-post*
 - the agent knows all agents' types.

Ex-interim expected utility

Definition (*Ex-interim* expected utility)

Agent i 's *ex-interim expected utility* in a Bayesian game (N, A, Θ, p, u) , where i 's type is θ_i and where the agents' strategies are given by the mixed strategy profile s , is defined as

$$EU_i(s|\theta_i) = \sum_{\theta_{-i} \in \Theta_{-i}} p(\theta_{-i}|\theta_i) \sum_{a \in A} \left(\prod_{j \in N} s_j(a_j|\theta_j) \right) u_i(a, \theta_{-i}, \theta_i).$$

- i must consider every θ_{-i} and every a in order to evaluate $u_i(a, \theta_i, \theta_{-i})$.
- i must weight this utility value by:
 - the probability that a would be realized given all players' mixed strategies and types;
 - the probability that the other players' types would be θ_{-i} given that his own type is θ_i .

Ex-ante expected utility

Definition (*Ex-ante* expected utility)

Agent i 's **ex-ante expected utility** in a Bayesian game (N, A, Θ, p, u) , where the agents' strategies are given by the mixed strategy profile s , is defined as

$$EU_i(s) = \sum_{\theta_i \in \Theta_i} p(\theta_i) EU_i(s|\theta_i)$$

or equivalently as

$$EU_i(s) = \sum_{\theta \in \Theta} p(\theta) \sum_{a \in A} \left(\prod_{j \in N} s_j(a_j|\theta_j) \right) u_i(a, \theta).$$

Ex-post expected utility

Definition (*Ex-post* expected utility)

Agent i 's **ex-post expected utility** in a Bayesian game (N, A, Θ, p, u) , where the agents' strategies are given by s and the agent' types are given by θ , is defined as

$$EU_i(s, \theta) = \sum_{a \in A} \left(\prod_{j \in N} s_j(a_j | \theta_j) \right) u_i(a, \theta).$$

- The only uncertainty here concerns the other agents' mixed strategies, since i knows everyone's type.

Best response

Definition (Best response in a Bayesian game)

The set of agent i 's **best responses** to mixed strategy profile s_{-i} are given by

$$BR_i(s_{-i}) = \arg \max_{s'_i \in S_i} EU_i(s'_i, s_{-i}).$$

- it may seem odd that BR is calculated based on i 's *ex-ante* expected utility.
- However, write $EU_i(s)$ as $\sum_{\theta_i \in \Theta_i} p(\theta_i) EU_i(s|\theta_i)$ and observe that $EU_i(s'_i, s_{-i}|\theta_i)$ does not depend on strategies that i would play if his type were not θ_i .
- Thus, we are in fact performing independent maximization of i 's *ex-interim* expected utility conditioned on each type that he could have.

Nash equilibrium

Definition (Bayes-Nash equilibrium)

A **Bayes-Nash equilibrium** is a mixed strategy profile s that satisfies $\forall i \ s_i \in BR_i(s_{-i})$.

- we can also construct an induced normal form for Bayesian games
- the numbers in the cells will correspond to *ex-ante* expected utilities
 - however as argued above, as long as the strategy space is unchanged, best responses don't change between the *ex-ante* and *ex-interim* cases.

ex-post Equilibrium

Definition (*ex-post* equilibrium)

A ***ex-post* equilibrium** is a mixed strategy profile s that satisfies $\forall \theta, \forall i, s_i \in \arg \max_{s'_i \in S_i} EU_i(s'_i, s_{-i}, \theta)$.

- somewhat similar to **dominant strategy**, but not quite
 - EP: agents do not need to have accurate beliefs about the type distribution
 - DS: agents do not need to have accurate beliefs about others' strategies