

# Computing Domination; Correlated Equilibria

## CPSC 532A Lecture 6

# Lecture Overview

- 1 Recap
- 2 Fun Game
- 3 Computational Problems Involving Domination
- 4 Rationalizability
- 5 Correlated Equilibrium

# Computing equilibria of zero-sum games

$$\begin{array}{ll}
 \text{minimize} & U_1^* \\
 \text{subject to} & \sum_{a_2 \in A_2} u_1(a_1, a_2) \cdot s_2^{a_2} \leq U_1^* \quad \forall a_1 \in A_1 \\
 & \sum_{a_2 \in A_2} s_2^{a_2} = 1 \\
 & s_2^{a_2} \geq 0 \quad \forall a_2 \in A_2
 \end{array}$$

- This formulation gives us the minmax strategy for player 2.
- To get the minmax strategy for player 1, we need to solve a second (analogous) LP.

# Computing Maxmin Strategies in General-Sum Games

To compute a maxmin strategy for player 1 in an arbitrary 2-player game  $G$ :

- Create a new game  $G'$  where player 2's payoffs are just the negatives of player 1's payoffs.
- By the minmax theorem, equilibrium strategies for player 1 in  $G'$  are equivalent to a maxmin strategies
- Thus, to find a maxmin strategy for  $G$ , find an equilibrium strategy for  $G'$ .

# Domination

- Let  $s_i$  and  $s'_i$  be two strategies for player  $i$ , and let  $S_{-i}$  be the set of all possible strategy profiles for the other players

## Definition

$s_i$  **strictly dominates**  $s'_i$  if  $\forall s_{-i} \in S_{-i}, u_i(s_i, s_{-i}) > u_i(s'_i, s_{-i})$

## Definition

$s_i$  **weakly dominates**  $s'_i$  if  $\forall s_{-i} \in S_{-i}, u_i(s_i, s_{-i}) \geq u_i(s'_i, s_{-i})$  and  $\exists s_{-i} \in S_{-i}, u_i(s_i, s_{-i}) > u_i(s'_i, s_{-i})$

## Definition

$s_i$  **very weakly dominates**  $s'_i$  if  $\forall s_{-i} \in S_{-i}, u_i(s_i, s_{-i}) \geq u_i(s'_i, s_{-i})$

# Iterated Removal of Dominated Strategies

- This process preserves Nash equilibria.
  - strict dominance: all equilibria preserved.
  - weak or very weak dominance: at least one equilibrium preserved.
- Thus, it can be used as a preprocessing step before computing an equilibrium
  - Some games are solvable using this technique.
- What about the order of removal when there are multiple dominated strategies?
  - strict dominance: doesn't matter.
  - weak or very weak dominance: can affect which equilibria are preserved.

# Lecture Overview

- 1 Recap
- 2 Fun Game**
- 3 Computational Problems Involving Domination
- 4 Rationalizability
- 5 Correlated Equilibrium

# Fun game

	$L$	$H$	$S$
$L$	90, 90	0, 0	0, 40
$B$	0, 0	180, 180	0, 40



# Fun game

	$L$	$H$	$S$
$L$	90, 90	0, 0	400, 40
$B$	0, 0	180, 180	0, 40

# Fun game

	$L$	$H$	$S$
$L$	90, 90	0, 0	0, 40; 400, 40
$B$	0, 0	180, 180	0, 40

- What's the equilibrium?

# Fun game

	$L$	$H$	$S$
$L$	90, 90	0, 0	0, 40; 400, 40
$B$	0, 0	180, 180	0, 40

- What's the equilibrium?
  - 50-50 L-H dominates  $S$  for column, so we have a standard coordination game.

# Fun game

	$L$	$H$	$S$
$L$	90, 90	0, 0	0, 40; 400, 40
$B$	0, 0	180, 180	0, 40

- What's the equilibrium?
  - 50-50 L-H dominates  $S$  for column, so we have a standard coordination game.
- What happens when people play?

# Fun game

	<i>L</i>	<i>H</i>	<i>S</i>
<i>L</i>	90, 90	0, 0	0, 40; 400, 40
<i>B</i>	0, 0	180, 180	0, 40

- What's the equilibrium?
  - 50-50 L-H dominates *S* for column, so we have a standard coordination game.
- What happens when people play?
  - with 0, 40, 96% row and 84% column choose the high payoff *H*, coordination occurs 80% of the time.
  - with 400, 40, 64% row and 76% column chose *H*; coordination on H,H 32% of the time, coordination on L,L 16% of the time, uncoordinated over half the time

# Lecture Overview

- 1 Recap
- 2 Fun Game
- 3 Computational Problems Involving Domination**
- 4 Rationalizability
- 5 Correlated Equilibrium

# Computational Problems in Domination

- Identifying strategies **dominated by a pure strategy**
- Identifying strategies **dominated by a mixed strategy**
- Identifying strategies **that survive iterated elimination**
- Asking whether a strategy survives iterated elimination under **all elimination orderings**
- We'll assume that  $i$ 's utility function is strictly positive everywhere (why is this OK?)

# Is $s_i$ strictly dominated by any pure strategy?

Try to identify some pure strategy that is strictly better than  $s_i$  for any pure strategy profile of the others.

```

for all pure strategies  $a_i \in A_i$  for player  $i$  where  $a_i \neq s_i$  do
   $dom \leftarrow true$ 
  for all pure strategy profiles  $a_{-i} \in A_{-i}$  for the players other than  $i$ 
  do
    if  $u_i(s_i, a_{-i}) \geq u_i(a_i, a_{-i})$  then
       $dom \leftarrow false$ 
      break
    end if
  end for
  if  $dom = true$  then return true
end for
return false

```



# Is $s_i$ strictly dominated by any pure strategy?

Try to identify some pure strategy that is strictly better than  $s_i$  for any pure strategy profile of the others.

```

for all pure strategies  $a_i \in A_i$  for player  $i$  where  $a_i \neq s_i$  do
   $dom \leftarrow true$ 
  for all pure strategy profiles  $a_{-i} \in A_{-i}$  for the players other than  $i$ 
  do
    if  $u_i(s_i, a_{-i}) \geq u_i(a_i, a_{-i})$  then
       $dom \leftarrow false$ 
      break
    end if
  end for
  if  $dom = true$  then return  $true$ 
end for
return  $false$ 

```

- What is the complexity of this procedure?
- Why don't we have to check mixed strategies of  $-i$ ?
- Minor changes needed to test for weak, very weak dominance.

# Constraints for determining whether $s_i$ is strictly dominated by any mixed strategy

$$\sum_{j \in A_i} p_j u_i(a_j, a_{-i}) > u_i(s_i, a_{-i}) \quad \forall a_{-i} \in A_{-i}$$

$$p_j \geq 0 \quad \forall j \in A_i$$

$$\sum_{j \in A_i} p_j = 1$$

# Constraints for determining whether $s_i$ is strictly dominated by any mixed strategy

$$\sum_{j \in A_i} p_j u_i(a_j, a_{-i}) > u_i(s_i, a_{-i}) \quad \forall a_{-i} \in A_{-i}$$

$$p_j \geq 0 \quad \forall j \in A_i$$

$$\sum_{j \in A_i} p_j = 1$$

- **What's wrong** with this program?

# Constraints for determining whether $s_i$ is strictly dominated by any mixed strategy

$$\sum_{j \in A_i} p_j u_i(a_j, a_{-i}) > u_i(s_i, a_{-i}) \quad \forall a_{-i} \in A_{-i}$$

$$p_j \geq 0 \quad \forall j \in A_i$$

$$\sum_{j \in A_i} p_j = 1$$

- **What's wrong** with this program?
  - **strict inequality** in the first constraint means we don't have an LP

# LP for determining whether $s_i$ is strictly dominated by any mixed strategy

$$\begin{array}{ll}
 \text{minimize} & \sum_{j \in A_i} p_j \\
 \text{subject to} & \sum_{j \in A_i} p_j u_i(a_j, a_{-i}) \geq u_i(s_i, a_{-i}) \quad \forall a_{-i} \in A_{-i} \\
 & p_j \geq 0 \quad \forall j \in A_i
 \end{array}$$

- This is clearly an LP. **Why is it a solution** to our problem?

# LP for determining whether $s_i$ is strictly dominated by any mixed strategy

$$\begin{array}{ll}
 \text{minimize} & \sum_{j \in A_i} p_j \\
 \text{subject to} & \sum_{j \in A_i} p_j u_i(a_j, a_{-i}) \geq u_i(s_i, a_{-i}) \quad \forall a_{-i} \in A_{-i} \\
 & p_j \geq 0 \quad \forall j \in A_i
 \end{array}$$

- This is clearly an LP. **Why is it a solution** to our problem?
  - if a solution exists with  $\sum_j p_j < 1$  then we can add  $1 - \sum_j p_j$  to some  $p_k$  and we'll have a dominating mixed strategy (since utility was assumed to be positive everywhere)
- Our original approach works for very weak domination
- For weak domination we can use that program with a different objective function trick.

# Identifying strategies that survive iterated elimination

- This can be done by repeatedly solving our LPs: solving a polynomial number of LPs is still in  $\mathcal{P}$ .
  - Checking whether every pure strategy of every player is dominated by any other mixed strategy requires us to solve at worst  $\sum_{i \in N} |A_i|$  linear programs.
  - Each step removes one pure strategy for one player, so there can be at most  $\sum_{i \in N} (|A_i| - 1)$  steps.
  - Thus we need to solve  $O((n \cdot \max_i |A_i|)^2)$  linear programs.

## Further questions about iterated elimination

- 1 **(Strategy Elimination)** Does there exist some elimination path under which the strategy  $s_i$  is eliminated?
- 2 **(Reduction Identity)** Given action subsets  $A'_i \subseteq A_i$  for each player  $i$ , does there exist a maximally reduced game where each player  $i$  has the actions  $A'_i$ ?
- 3 **(Uniqueness)** Does every elimination path lead to the same reduced game?
- 4 **(Reduction Size)** Given constants  $k_i$  for each player  $i$ , does there exist a maximally reduced game where each player  $i$  has exactly  $k_i$  actions?



## Further questions about iterated elimination

- 1 **(Strategy Elimination)** Does there exist some elimination path under which the strategy  $s_i$  is eliminated?
  - 2 **(Reduction Identity)** Given action subsets  $A'_i \subseteq A_i$  for each player  $i$ , does there exist a maximally reduced game where each player  $i$  has the actions  $A'_i$ ?
  - 3 **(Uniqueness)** Does every elimination path lead to the same reduced game?
  - 4 **(Reduction Size)** Given constants  $k_i$  for each player  $i$ , does there exist a maximally reduced game where each player  $i$  has exactly  $k_i$  actions?
- For **iterated strict dominance** these problems are all in  $\mathcal{P}$ .
  - For **iterated weak or very weak dominance** these problems are all  $\mathcal{NP}$ -complete.

# Lecture Overview

- 1 Recap
- 2 Fun Game
- 3 Computational Problems Involving Domination
- 4 Rationalizability**
- 5 Correlated Equilibrium

# Rationalizability

- Rather than ask what is irrational, ask what is a best response to **some** beliefs about the opponent
  - assumes opponent is rational
  - assumes opponent knows that you and the others are rational
  - ...
- Examples
  - is *heads* rational in matching pennies?

# Rationalizability

- Rather than ask what is irrational, ask what is a best response to **some** beliefs about the opponent
  - assumes opponent is rational
  - assumes opponent knows that you and the others are rational
  - ...
- Examples
  - is *heads* rational in matching pennies?
  - is *cooperate* rational in prisoner's dilemma?

# Rationalizability

- Rather than ask what is irrational, ask what is a best response to **some** beliefs about the opponent
  - assumes opponent is rational
  - assumes opponent knows that you and the others are rational
  - ...
- Examples
  - is *heads* rational in matching pennies?
  - is *cooperate* rational in prisoner's dilemma?
- Will there always exist a rationalizable strategy?

# Rationalizability

- Rather than ask what is irrational, ask what is a best response to **some** beliefs about the opponent
  - assumes opponent is rational
  - assumes opponent knows that you and the others are rational
  - ...
- Examples
  - is *heads* rational in matching pennies?
  - is *cooperate* rational in prisoner's dilemma?
- Will there always exist a rationalizable strategy?
  - Yes, equilibrium strategies are always rationalizable.

# Rationalizability

- Rather than ask what is irrational, ask what is a best response to **some** beliefs about the opponent
  - assumes opponent is rational
  - assumes opponent knows that you and the others are rational
  - ...
- Examples
  - is *heads* rational in matching pennies?
  - is *cooperate* rational in prisoner's dilemma?
- Will there always exist a rationalizable strategy?
  - Yes, equilibrium strategies are always rationalizable.
- Furthermore, in two-player games, rationalizable  $\Leftrightarrow$  survives iterated removal of strictly dominated strategies.

# Lecture Overview

- 1 Recap
- 2 Fun Game
- 3 Computational Problems Involving Domination
- 4 Rationalizability
- 5 Correlated Equilibrium**



# Pithy Quote

*If there is intelligent life on other planets, in a majority of them, they would have discovered correlated equilibrium before Nash equilibrium.*

– Roger Myerson

# Examples

- Consider again Battle of the Sexes.
  - Intuitively, the best outcome seems a 50-50 split between  $(F, F)$  and  $(B, B)$ .
  - But there's no way to achieve this, so either someone loses out (unfair) or both players often miscoordinate
- Another classic example: traffic game

	<i>go</i>	<i>wait</i>
<i>go</i>	-100, -100	10, 0
<i>B</i>	0, 10	-10, -10

# Intuition

- What is the natural solution here?

# Intuition

- What is the natural solution here?
  - A traffic light: a fair randomizing device that tells one of the agents to go and the other to wait.
- Benefits:
  - the negative payoff outcomes are completely avoided
  - fairness is achieved
  - the sum of social welfare exceeds that of any Nash equilibrium
- We could use the same idea to achieve the fair outcome in battle of the sexes.
- Our example presumed that everyone perfectly observes the random event; not required.
- More generally, some random variable with a commonly known distribution, and a private signal to each player about the outcome.
  - signal doesn't determine the outcome or others' signals; however, correlated

# Formal definition

## Definition (Correlated equilibrium)

Given an  $n$ -agent game  $G = (N, A, u)$ , a **correlated equilibrium** is a tuple  $(v, \pi, \sigma)$ , where  $v$  is a tuple of random variables  $v = (v_1, \dots, v_n)$  with respective domains  $D = (D_1, \dots, D_n)$ ,  $\pi$  is a joint distribution over  $v$ ,  $\sigma = (\sigma_1, \dots, \sigma_n)$  is a vector of mappings  $\sigma_i : D_i \mapsto A_i$ , and for each agent  $i$  and every mapping  $\sigma'_i : D_i \mapsto A_i$  it is the case that

$$\sum_{d \in D} \pi(d) u_i(\sigma_1(d_1), \dots, \sigma_n(d_n)) \geq \sum_{d \in D} \pi(d) u_i(\sigma'_1(d_1), \dots, \sigma'_n(d_n))$$

# Existence

## Theorem

For every Nash equilibrium  $\sigma^*$  there exists a *corresponding correlated equilibrium*  $\sigma$ .

- This is easy to show:
  - let  $D_i = A_i$
  - let  $\pi(d) = \prod_{i \in N} \sigma_i^*(d_i)$
  - $\sigma_i$  maps each  $d_i$  to the corresponding  $a_i$ .
- Thus, correlated equilibria always exist

# Remarks

- Not every correlated equilibrium is equivalent to a Nash equilibrium
  - thus, correlated equilibrium is a **weaker notion** than Nash
- Any **convex combination of the payoffs** achievable under correlated equilibria is itself realizable under a correlated equilibrium
  - start with the Nash equilibria (each of which is a CE)
  - introduce a second randomizing device that selects which CE the agents will play
  - regardless of the probabilities, no agent has incentive to deviate
  - the probabilities can be adjusted to achieve any convex combination of the equilibrium payoffs
  - the randomizing devices can be combined