

# FAST MARCHING METHODS FOR STATIONARY HAMILTON-JACOBI EQUATIONS WITH AXIS-ALIGNED ANISOTROPY

KEN ALTON AND IAN M. MITCHELL\*

**Date.** August 8, 2008

**Technical Report Number.** TR-2008-02

**Accepted for Publication in.** SIAM Journal on Numerical Analysis.

**Abstract.** The Fast Marching Method (FMM) has proved to be a very efficient algorithm for solving the isotropic Eikonal equation. Because it is a minor modification of Dijkstra's algorithm for finding the shortest path through a discrete graph, FMM is also easy to implement. In this paper we describe a new class of Hamilton-Jacobi (HJ) PDEs with axis-aligned anisotropy which satisfy a causality condition for standard finite difference schemes on orthogonal grids and can hence be solved using the FMM; the only modification required to the algorithm is in the local update equation for a node. This class of HJ PDEs has applications in anelliptic wave propagation and robotic path planning, and brief examples are included. Since our class of HJ PDEs and grids permit asymmetries, we also examine some methods of improving the efficiency of the local update that do not require symmetric grids and PDEs. Finally, we include explicit update formulas for variations of the Eikonal equation that use the Manhattan, Euclidean and infinity norms on orthogonal grids of arbitrary dimension and with variable node spacing.

**Key words.** fast marching method, anisotropic optimal control, Hamilton-Jacobi equation, viscosity solution

**AMS subject classifications.** 35F30, 49L20, 49L25, 49N90, 65N06, 65N12

This technical report is an extended version of a paper that has been accepted for publication in SIAM Journal on Numerical Analysis [3]. This version contains several proofs that were omitted from the journal submission, including a proof that the output of FMM is the solution to the discretized HJ PDE and a proof that the solution to the discretized equation converges to the viscosity solution of the HJ PDE as the grid spacing approaches zero.

This technical report supercedes [2] in nearly all sections. One important exception is that [2] examines a dual relationship concerning the action set of a control-theoretic formulation (1.2) and the  $G$  function in norm-like formulation (2.2). This examination has been left out here.

**1. Introduction.** The Fast Marching Method (FMM) [31, 22] has become a popular algorithm to use when solving the Dirichlet problem for an isotropic static Hamilton-Jacobi Partial Differential Equation (HJ PDE), also known as the Eikonal equation  $\|Du(x)\|_2 = c(x)$ . FMM has proven to be particularly efficient in practice because it can approximately solve this problem in a single pass through the nodes of a grid. It is also straightforward to implement, requiring only a small modification of Dijkstra's algorithm [10], which is a popular method for finding the shortest path through a graph.

While the isotropic case is the most common, there are applications which require solution of anisotropic HJ PDEs. Unfortunately, FMM produces a correct approxima-

---

\*Department of Computer Science, University of British Columbia, Vancouver, BC V6T 1Z4, Canada ({kalton, mitchell}@cs.ubc.ca, <http://www.cs.ubc.ca/~mitchell>). This work was supported by a grant from the National Science and Engineering Research Council of Canada.

tion only under certain causality conditions on the values of nodes and their neighbors. This limitation has motivated the development of a more generally applicable version of FMM called Ordered Upwind Methods (OUMs) [27] and also several recent works such as [33, 15, 21] on sweeping methods. However, OUMs are much more complex to implement than FMM, and sweeping methods can be much less efficient for problems with curved characteristics and practical grid sizes [14, 13].

Consequently, we have motivation to seek classes of anisotropic problems to which FMM might still be applied. One such class of problems was identified in [26] and includes the Eikonal equation where an energy norm replaces the standard Euclidean norm. In [1] we identified another such class of problems. Because its characteristics are minimum time paths to the boundary, the Eikonal equation has often been proposed for robotic path planning; for example, see [17]. However, for some robots using the Euclidean norm in this equation is inappropriate. Consider a robot arm, where each joint has its own motor. If each motor can rotate at some maximum speed independent of the action of the other motors, then the action of the whole arm is best bounded in an appropriately-scaled infinity norm. The corresponding Eikonal equation should use the dual Manhattan norm and is thus anisotropic. Other scenarios where such problems arise were considered in [1]—such as planning collision free optimal paths for multiple robots—and experimental evidence suggested that FMM would be successful on these problems.

As a group, the anisotropy in these problems is axis-aligned. In this paper we describe a broader class of such axis-aligned problems (Section 2) and demonstrate that FMM can be applied to approximate their solution on axis-aligned orthogonal grids without modification of the algorithm beyond the local update function for a single node (Section 3). The examples (Section 5) include an anelliptic wave propagation problem and a new multirobot scenario. In Appendix A, we propose some methods by which the local update’s efficiency might be improved even if the grid and/or PDE lack symmetry. Lastly, in Appendix B, we provide analytic update formulas for the Eikonal equation with the  $p = 1, 2,$  and  $\infty$  norms on variably-spaced orthogonal grids in any dimension.

**1.1. The Problem.** The Dirichlet problem of a static HJ PDE is to find a function  $u$ , such that

$$H(x, Du(x)) = 0, \quad x \in \Omega \tag{1.1a}$$

$$u(x) = g(x), \quad x \in \partial\Omega, \tag{1.1b}$$

where  $Du(x)$  is the gradient of  $u$  at  $x$ ,  $\Omega \subset \mathbb{R}^d$  is a bounded Lipschitz domain, and  $\partial\Omega$  is the domain’s boundary. In general, it is not possible to find a classical solution to the Dirichlet problem (1.1) where  $u$  is differentiable for all  $x$ , so we seek instead the viscosity solution [8], a unique weak solution which is continuous and almost everywhere differentiable.

To appreciate the difference between isotropic and anisotropic problems it is useful to consider a control-theoretic formulation of the Hamiltonian,

$$H(x, q) = \max_{a \in \mathcal{A}(x)} (-q \cdot a) - 1, \tag{1.2}$$

where  $a$  is an action and  $\mathcal{A}(x) \subset \mathbb{R}^d$  is a compact, convex action set containing the origin in its interior. In an isotropic problem  $\mathcal{A}(x)$  is a hypersphere centered on the origin for all  $x$ , although its radius may depend on  $x$ . In such a problem (1.2) reduces




FIG. 1.1. Orthogonal grids combining discretizations  $\underline{\Omega}$  and  $\partial\underline{\Omega}$ . (a) boundary conditions are given around the outside of  $\Omega$ . (b) boundary conditions are given on the inside of  $\Omega$ .

to

$$H(x, q) = \|q\|_2 - c(x), \quad (1.3)$$

where  $c(x) = 1/r(x)$  and  $r(x)$  is the radius of the hyperspherical  $\mathcal{A}(x)$ . In this case (1.1a) becomes the Eikonal equation. For an anisotropic problem  $\mathcal{A}(x)$  is not always a origin-centered hypersphere. Since not all Hamiltonians  $H$  fit the control-theoretic formulation, more generally, for an isotropic problem the set of  $q$  solving  $H(x, q) = 0$  is the surface of a origin-centered hypersphere. Several examples of anisotropic problems that do not fit this criterion are included in Sections 2.2 and 5.

**1.2. The Fast Marching Method.** Since we typically cannot solve for the viscosity solution exactly, we compute an approximate solution  $\underline{u}$  on an axis-aligned orthogonal grid with nodes forming both a discretization  $\underline{\Omega}$  of  $\Omega$ , and a discretization  $\partial\underline{\Omega}$  of  $\partial\Omega$ ; for example, see Figure 1.2. We take  $\underline{\Omega}$  and  $\partial\underline{\Omega}$  to be disjoint sets. We allow any axis-aligned orthogonal grid, including those with node spacing that varies between dimensions and within a single dimension; the latter capability makes it easier to more accurately manage an irregular boundary [13]. It is important that the orthogonal grid and the Hamiltonian  $H$  are aligned to the same axis. What it means for  $H$  to be aligned to an axis is explained in section 2.

Let  $\mathcal{N}(\underline{x})$  be the set of neighbors of node  $\underline{x} \in \underline{\Omega}$ . Whenever we refer to a simplex of a node  $\underline{x}$ , we mean a simplex specified by the node  $\underline{x}$  and  $d$  of its neighbors, each in a distinct dimension. Since we are restricted to orthogonal grids, each simplex of  $\underline{x}$  corresponds to a particular orthant.

Informally, we refer to  $\underline{u}(\underline{x})$  as the value of node  $\underline{x}$ . In the sequel, we may use  $\underline{u}$  to refer to either the values of the nodes in the operation or output of FMM or to the solution of the discretized PDE (3.3). This ambiguity becomes less bothersome when we point out in Proposition 3.2 that the output of FMM is in fact the solution to (3.3).

Algorithm 1 outlines a simple dynamic programming algorithm. The algorithm can become either Dijkstra's algorithm or FMM depending on the choice of the `Update` function. Consider, for example, the `Update` function in the context of optimal control, where we are computing the minimal cost over all possible paths. For Dijkstra's algorithm, `Update` computes  $\underline{u}(\underline{x}_0)$  as a simple minimization over the neighboring

nodes of  $\underline{x}_0$  of the path costs to  $\underline{x}_0$  via each neighbor. For FMM, the **Update** function computes  $u(\underline{x}_0)$  as a minimization over the neighboring simplices of  $\underline{x}_0$  of the minimum path costs to  $\underline{x}_0$  via each simplex.

The **Update** function must satisfy a *causality* property in order for Algorithm 1 to terminate with a correct solution: **Update** must compute a node value  $\underline{u}(\underline{x})$  based only on information from neighboring nodes with smaller values, so that  $\underline{u}$  is computed in increasing order of  $\underline{u}(\underline{x})$  [30, 23]. In Dijkstra’s algorithm and FMM for a standard Euclidean norm Eikonal equation on an orthogonal grid, this property is automatic. A major contribution of this paper is to demonstrate that for a class of static HJ PDEs with axis-aligned anisotropy, an **Update** function that is consistent with the PDE and satisfies the causality property can be defined; and thus, FMM can be used.

```

1 foreach  $\underline{x} \in \underline{\Omega}$  do  $u(\underline{x}) \leftarrow \infty$ 
2 foreach  $\underline{x} \in \partial\underline{\Omega}$  do  $u(\underline{x}) \leftarrow g(\underline{x})$ 
3  $\mathcal{Q} \leftarrow \underline{\Omega} \cup \partial\underline{\Omega}$ 
4 while  $\mathcal{Q} \neq \emptyset$  do
5    $\underline{y} \leftarrow \text{ExtractMin}(\mathcal{Q})$ 
6   foreach  $\underline{x}_0 \in (\mathcal{N}(\underline{y}) \cap \mathcal{Q}) \setminus \partial\underline{\Omega}$  do  $u(\underline{x}_0) \leftarrow \text{Update}(\underline{x}_0, \underline{u})$ 
7 end

```

**Algorithm 1:** Dynamic Programming Algorithm.

While the **Update** function in Algorithm 1 is determined by the underlying equation which we seek to solve, it is assumed that its execution time is independent of grid resolution and hence it does not affect the algorithm’s asymptotic complexity. The **Update** functions in this paper maintain this property. FMM is usually described as being  $\mathcal{O}(n \log n)$ , where  $n = |\underline{\Omega}|$  is the number of grid points in the discretized domain. This complexity is derived by noting that each node is removed from  $\mathcal{Q}$  once by **ExtractMin** and, in the usual binary heap implementation of  $\mathcal{Q}$ , extraction of the minimum value node costs  $\mathcal{O}(\log |\mathcal{Q}|) \leq \mathcal{O}(\log n)$ . Note that the heap need only sort nodes with finite values. Because we restrict our modifications of Algorithm 1 to the **Update** function, all of the results here can be used with other versions of FMM; for example, the  $\mathcal{O}(n)$  algorithm described in [32], which uses an untidy priority queue for  $\mathcal{Q}$  to reduce the cost of **ExtractMin** and hence the whole algorithm. However, for implementation simplicity we have used the standard binary heap version of  $\mathcal{Q}$  in our experiments.

**1.3. Related Work.** The first Dijkstra-like method for a first-order semi-Lagrangian discretization of the isotropic Eikonal PDE on an orthogonal grid was developed in [30]. The Dijkstra-like FMM was later independently developed in [22] for the first-order upwind Eulerian finite-difference discretization of the same Eikonal PDE. FMM was then extended to handle higher-order upwind discretizations on grids and unstructured meshes in  $\mathbb{R}^n$  and on manifolds [16, 24, 26]. In [23] it was shown that Dijkstra’s method on a uniform orthogonal grid produces the solution for the anisotropic maximum norm Eikonal equation. By solving an isotropic problem on a manifold and then projecting the solution into a subspace, FMM can solve certain anisotropic problems [26]; for example, (1.2) with a constant elliptic  $\mathcal{A}(x) = \mathcal{A}$  can be solved by running isotropic FMM on an appropriately tilted planar manifold and then projecting away one dimension. Some anisotropic etching problems have also been solved using FMM [19].

The fact that correct operation of Dijkstra-like algorithms for approximating the Eikonal PDE requires the causality property that  $\underline{u}(\underline{x})$  can be written only in terms of smaller values  $\underline{u}$  at neighboring nodes was stated in [30], but a reader might incorrectly infer from further comments in that paper that such algorithms would not work for any unstructured grid or anisotropic problem. That FMM is applicable for any consistent, orthogonal, causality satisfying, finite-difference discretization of a general static convex HJ PDE is stated in [23]; however, it is now understood that this criterion applies even more generally, since a Dijkstra-like method can be used to efficiently solve on a graph any nonlinear system of equations for which  $\underline{u}(\underline{x})$  is dependent only on smaller values  $\underline{u}$  at neighboring nodes. A sufficient criterion (see Section 2.1) under which FMM can be used for orthogonal, finite-difference discretizations of static HJ PDEs—now commonly referred to as “Osher’s criterion”—is widely attributed to an unpublished work by Osher & Helmsen, but the earliest published description seems to be [19]. While it is stronger than the causality conditions described earlier, it is useful because it is stated as a condition on the analytic Hamiltonian instead of the equations created by the discretization. In this paper we likewise seek conditions under which FMM is applicable that are closer to the problem’s definition than the algorithm’s implementation.

OUMs [27, 28] can solve general convex anisotropic problems on unstructured grids with an asymptotic complexity only a constant factor (related to the degree of anisotropy) worse than FMM. FMM fails for these general problems because the neighboring simplex from which the characteristic approaches a node  $\underline{x}_0$  may contain another node  $\underline{x}$  such that causality does not hold:  $\underline{u}(\underline{x}_0) < \underline{u}(\underline{x})$ . OUM avoids this difficulty by searching along the active front to find a set of neighboring nodes (which may not be direct neighbors of  $\underline{x}_0$ ) whose values have been accepted, and then constructing a virtual simplex with these nodes from which to update  $\underline{u}(\underline{x}_0)$ . Although this search along the active front does not degrade the asymptotic complexity, it does significantly increase the computational cost in practice. This effect can be partially mitigated by using nontrivial data structures such as  $2^d$ -trees to speed up the search.

An alternative to these single pass (or label setting) algorithms are the sweeping (or label correcting) algorithms, which are often even simpler to implement than FMM. Sweeping algorithms are also capable of handling anisotropic and even non-convex problems. The simplest sweeping algorithm is to just iterate through the grid updating each node in a Gauss-Seidel (GS) fashion (so a new value for a node is used immediately in subsequent updates) until  $\underline{u}$  converges. GS converges quickly if the node update order is aligned with the characteristics of the solution, so better sweeping algorithms [9, 7, 33, 15, 21] alternate among a collection of static node orderings so that all possible characteristic directions will align with at least one ordering. It is argued in [33] that these methods achieve  $\mathcal{O}(n)$  asymptotic complexity (assuming that the node orderings are already determined); however, unlike FMM and OUM the constant depends on the problem. For practical grid resolutions on problems with curved characteristics FMM does better despite the difference in asymptotic complexity [14, 13].

There are also a number of sweeping algorithms which use dynamic node orderings; for example [20, 6]. These algorithms attempt to approximate the optimal ordering generated by single-pass methods such as FMM without the overhead associated with managing an accurate queue. These methods have been demonstrated to be comparable to or better than single-pass methods for certain problems and grid resolutions [20, 6]. However, in general these methods may need to revisit nodes

multiple times.

Accurate robotic path planning is only required in cluttered environments where optimal paths—and hence the characteristics of the HJ PDE—are not straight. No alternative algorithm proposed approaches the simple implementation and guaranteed speed of FMM for these types of problems. Consequently, we set out in this paper to characterize another class of anisotropic HJ PDEs for which FMM will work, and also to explore their efficient implementation. It should be noted that the update procedures discussed in this paper can be applied to any of the sweeping algorithms without modification.

**2. Class of Hamiltonians.** FMM can be extended to handle a class of axis-aligned anisotropic problems, defined by a restriction of the Hamiltonian  $H$  to that satisfying Properties 1 to 4. We let  $q, \tilde{q} \in \mathbb{R}^d$  and make the definitions:

DEFINITION 2.1. Write  $q \succeq \tilde{q}$  if  $q_j \tilde{q}_j \geq 0$  and  $|q_j| \geq |\tilde{q}_j|$ , for  $1 \leq j \leq d$ .

DEFINITION 2.2. Write  $q \succ \tilde{q}$  if (i)  $q \neq 0$  and (ii)  $q_j \tilde{q}_j \geq 0$  and  $|q_j| > |\tilde{q}_j|$  or  $q_j = \tilde{q}_j = 0$ , for  $1 \leq j \leq d$ .

The following properties are satisfied by  $H$ :

PROPERTY 1.  $H$  is continuous:  $H \in C(\Omega \times \mathbb{R}^d)$ .

PROPERTY 2.  $H$  is coercive:  $H(x, q) \rightarrow \infty$  as  $\|q\| \rightarrow \infty$ , for all  $x \in \Omega$ .

PROPERTY 3.  $H$  is strictly compatible:  $H(x, 0) < 0$ , for all  $x \in \Omega$ .

PROPERTY 4.  $H$  is strictly one-sided monotone: If  $q \succ \tilde{q}$ , then  $H(x, q) > H(x, \tilde{q})$ .

We note that Properties 1, 2, and 3 are similar to some properties on the Hamiltonian in [6]. In this paper, we typically deal only with the `Update` function. For this reason, we usually consider a fixed  $x \in \Omega$  and may write  $H(q) = H(x, q)$  wherever no ambiguity results. When discussing properties of  $H$  these are in reference to the  $q$  parameter. The source of the *axis-aligned* description of the problem class is the strict one-sided monotonicity property of  $H$ .

**2.1. Connection to Osher’s criterion.** Although there are earlier statements of the conditions on node values under which a Dijkstra-like algorithm can or cannot be used to solve the problem [30, 22], in this section we outline the connection between the properties described above and Osher’s criterion [19] because the latter directly provides a condition on the Hamiltonian rather than on the solution values. In Section 3.3, we make the connection between Properties 1 to 4 and the earlier conditions.

Osher’s fast marching criterion is defined in [19, 29] as

$$q_j \frac{\partial H(x, q)}{\partial q_j} \geq 0$$

for  $1 \leq j \leq d$ . The authors state there that as long as this criterion is satisfied, a simple fast marching algorithm based on a one-sided upwind finite-difference discretization can be applied to solve the problem. However, we use Properties 1 to 4 instead of Osher’s criterion because Osher’s criterion requires  $H$  to be differentiable so that  $D_q H(x, q)$  exists, but we are interested in potentially nondifferentiable  $H$  (e.g., see Section 2.2). Note that strict one-sided monotonicity is applicable even when  $D_q H(x, q)$  does not exist for all  $x$ .

Propositions 2.3, 2.4, and 2.5 explain the relationship between strict one-sided monotonicity of  $H$  (Property 4) and Osher’s criterion. Proposition 2.3 shows that Property 4 implies one-sided monotonicity (Property 5). Then, Proposition 2.4 shows

that Property 5 is the same as Osher's criterion as long as  $H$  is differentiable. Finally, Proposition 2.5 demonstrates that Property 5 with the addition of one-sided homogeneity (Property 6) implies Property 4.

PROPERTY 5.  $H$  is one-sided monotone: If  $q \succeq \tilde{q}$ , then  $H(x, q) \geq H(x, \tilde{q})$ .

PROPOSITION 2.3. Let  $H$  be continuous (Property 1). Then strict one-sided monotonicity of  $H$  (Property 4) implies one-sided monotonicity of  $H$  (Property 5).

*Proof.* Let  $H$  be strictly one-sided monotone. Let  $q, \tilde{q} \in \mathbb{R}^d$  be such that  $q \succeq \tilde{q}$ . Let  $r \in \{-1, 1\}^d$  be such that

$$r_j = \begin{cases} +1, & \text{if } q_j \geq 0 \\ -1, & \text{otherwise} \end{cases}$$

and let  $\epsilon > 0$ . Note that  $q + \epsilon r \succ \tilde{q}$  and thus we have  $H(q + \epsilon r) > H(\tilde{q})$ . By the continuity of  $H$  we have

$$\lim_{\epsilon \rightarrow 0^+} H(q + \epsilon r) \geq H(\tilde{q})$$

and also

$$\lim_{\epsilon \rightarrow 0^+} H(q + \epsilon r) = H(q).$$

Therefore,  $H(q) \geq H(\tilde{q})$ .  $\square$

PROPOSITION 2.4. Let  $H$  be continuous (Property 1) and let  $D_q H(q)$  exist for all  $q \in \mathbb{R}^d$ . Then the following conditions on  $H$  are equivalent.

(a)  $q_j \frac{\partial H(q)}{\partial q_j} \geq 0$ , for all  $j$  and  $q \in \mathbb{R}^d$ .

(b)  $H$  is one-sided monotone (Property 5).

*Proof.* We begin by proving that (a) implies (b). Let  $q, \tilde{q} \in \mathbb{R}^d$  be such that  $q \succeq \tilde{q}$ . If  $q = \tilde{q}$  then  $H(q) = H(\tilde{q})$ . Otherwise, define the function  $\bar{q} : [0, 1] \rightarrow \mathbb{R}^d$  such that  $\bar{q}(t) = \tilde{q} + t(q - \tilde{q})$  to represent the line segment between  $\tilde{q}$  and  $q$  parameterized by  $t \in [0, 1]$ . Because  $q \succeq \tilde{q}$  we have

$$(q - \tilde{q})_j \bar{q}_j(t) \geq 0$$

for  $1 \leq j \leq d$  and for  $t \in [0, 1]$ . Thus, by condition (a) we have

$$(q - \tilde{q})_j \frac{\partial H(\bar{q}_j(t))}{\partial \bar{q}_j(t)} \geq 0, \quad (2.1)$$

for  $1 \leq j \leq d$  and for  $t \in [0, 1]$ . We know that

$$\begin{aligned} H(q) &= H(\tilde{q}) + \int_0^1 \frac{d\bar{q}(t)}{dt} \cdot D_q H(\bar{q}(t)) dt \\ &= H(\tilde{q}) + \int_0^1 (q - \tilde{q}) \cdot D_q H(\bar{q}(t)) dt \\ &= H(\tilde{q}) + \int_0^1 \sum_{i=1}^n (q - \tilde{q})_i \frac{\partial H(\bar{q}_i(t))}{\partial \bar{q}_i(t)} dt \\ &\geq H(\tilde{q}). \end{aligned}$$

The first equality follows from integrating the change in  $H$  along the line segment connecting  $\tilde{q}$  and  $q$ . The second equality is because the derivative  $\frac{d\bar{q}(t)}{dt}$  is simply

the vector  $q - \tilde{q}$ . The third equality breaks up the vector dot product into a sum of scalar products. The inequality results from (2.1) and the fact that an integral of a nonnegative function is nonnegative. Thus for all  $q, \tilde{q}$  such that  $q \triangleright \tilde{q}$ , including  $q = \tilde{q}$ , we have  $H(q) \geq H(\tilde{q})$ .

We now prove that (b) implies (a). Let  $q \in \mathbb{R}^d$  and  $1 \leq j \leq d$ . Define the function  $s : \mathbb{R} \rightarrow \{-1, +1\}$  such that

$$s(y) = \begin{cases} +1, & \text{if } y \geq 0 \\ -1, & \text{otherwise,} \end{cases}$$

let  $\epsilon > 0$ , and let  $e_j$  be the  $j$ th vector in the standard basis. Note that  $q + \epsilon s(q_j)e_j \triangleright q$  and thus by (b) we have  $H(q + \epsilon s(q_j)e_j) - H(q) \geq 0$ . Consequently, by the existence of  $D_q H(q)$  for all  $q \in \mathbb{R}^d$  we have

$$q_j \frac{\partial H(q)}{\partial q_j} = \lim_{\epsilon \rightarrow 0^+} q_j \frac{H(q + \epsilon s(q_j)e_j) - H(q)}{\epsilon s(q_j)} \geq 0.$$

□

The following property is used to state Proposition 2.5.

PROPERTY 6. *H is one-sided homogeneous:  $H(tq) - H(0) = t(H(q) - H(0))$  for all  $t \geq 0$  and  $q \in \mathbb{R}^d$ .*

PROPOSITION 2.5. *Let H satisfy Properties 1, 2, and 3, and let H be one-sided monotone (Property 5) and one-sided homogeneous (Property 6). Then H is strictly one-sided monotone (Property 4).*

*Proof.* Let  $q \triangleright \tilde{q}$ . Then  $q \triangleright \tilde{q}$  and  $H(q) \geq H(\tilde{q})$  by one-sided monotonicity.

First consider the case  $\tilde{q} = 0$ . Assume  $H(q) = H(\tilde{q}) = H(0)$ . By the one-sided homogeneity of  $H$ ,

$$\lim_{t \rightarrow \infty} [H(tq) - H(0)] = \lim_{t \rightarrow \infty} [t(H(q) - H(0))] = 0.$$

But by the coercivity of  $H$

$$\lim_{t \rightarrow \infty} [H(tq) - H(0)] = \infty,$$

since  $\lim_{t \rightarrow \infty} \|tq\| = \infty$  and by compatability  $H(0) < 0$ . Thus we have a contradiction and it must be that  $H(q) > H(\tilde{q})$ .

Second, consider the case where  $\tilde{q} \neq 0$ . Let  $\mathcal{J} = \{j \mid |q_j| > |\tilde{q}_j|\}$ . Note that by Definition 2.2 since  $\tilde{q} \neq 0$ , we have  $\mathcal{J} \neq \emptyset$  and  $\exists j \in \mathcal{J}$  such that  $\tilde{q}_j \neq 0$ . Define a scalar multiple of  $q$ :

$$\check{q} = tq = \left( \max_{j \in \mathcal{J}} \frac{|\tilde{q}_j|}{|q_j|} \right) q.$$

Since  $|q_j| > |\tilde{q}_j|, \forall j \in \mathcal{J}$ , we have  $0 < t < 1$ . Furthermore, for  $j \in \mathcal{J}$ ,

$$|\check{q}_j| = \left( \max_{j \in \mathcal{J}} \frac{|\tilde{q}_j|}{|q_j|} \right) |q_j| \geq |\tilde{q}_j|,$$

while for  $j \notin \mathcal{J}$ ,

$$\check{q}_j = tq_j = 0 = \tilde{q}_j.$$






FIG. 2.1. Contour plots of  $\|q\|_p$ .

Consequently,  $|\check{q}_j| \geq |\tilde{q}_j|$  for  $1 \leq j \leq d$ . Also, since  $t > 0$ , we have  $\check{q}_j \tilde{q}_j = tq_j \tilde{q}_j \geq 0$  for  $1 \leq j \leq d$ . This implies, by one-sided monotonicity of  $H$ , that  $H(\check{q}) \geq H(\tilde{q})$ . Moreover, by one-sided homogeneity of  $H$ ,  $H(\tilde{q}) - H(0) = H(tq) - H(0) = t(H(q) - H(0))$ . It follows that  $H(q) - H(0) = (H(\check{q}) - H(0))/t > H(\tilde{q}) - H(0)$ , since  $0 < t < 1$  and  $H(\check{q}) \geq H(0)$  by one-sided monotonicity. Therefore,  $H(q) > H(\check{q}) \geq H(\tilde{q})$ .  $\square$

We impose strict one-sided monotonicity on  $H$  because it guarantees a unique solution to a first-order upwind finite-difference discretization of (1.1a), as shown in Section 3.1. Simply imposing one-sided monotonicity on  $H$  or Osher's condition on differentiable  $H$  is not sufficient for a unique solution. However, Proposition 2.5 states that when  $H$  satisfies one-sided homogeneity in addition to one-sided monotonicity then it also satisfies strict one-sided monotonicity and there is a unique solution to the discretization. Moreover, by Propositions 2.4 and 2.5, when differentiable  $H$  satisfies one-sided homogeneity in addition to Osher's criterion then  $H$  also satisfies strict one-sided monotonicity and there is a unique solution to the discretization. Note that there exist conditions other than one-sided homogeneity, such as strict convexity, that in combination with Osher's criterion result in strict one-sided monotonicity of  $H$ .

**2.2. Example  $H$  Functions.** A Hamiltonian  $H$  that satisfies Properties 1 to 4 encompasses a fairly broad range of anisotropic problems. We consider examples of  $H$  that satisfy Properties 1 to 4. In particular, we look at the case

$$H(x, q) = G(x, q) - c(x), \quad (2.2)$$

where  $G$  is a  $p$ -norm or some variant and  $c$  is a positive cost. We must ensure that  $G$  is strictly one-sided monotone, which is not true of all norms.

The  $p$ -norm is a useful category of strictly one-sided monotone norms. Let a  $p$ -norm,  $\|\cdot\|_p$ , be defined by

$$\|q\|_p = \left( \sum_{j=1}^d |q_j|^p \right)^{1/p},$$

where  $p \geq 1$ . Commonly used  $p$ -norms, illustrated in Figure 2.1, are the Manhattan norm ( $p = 1$ ), the Euclidean norm ( $p = 2$ ), and the maximum norm ( $p = \infty$ ).

PROPOSITION 2.6.  $\|\cdot\|_p$  is strictly one-sided monotone, for  $p \geq 1$ .

*Proof.* Let  $q \succ \tilde{q}$ . We know that  $|q_j| \geq |\tilde{q}_j| \geq 0$  for all  $j$ , such that  $1 \leq j \leq d$ . Furthermore,  $|q_j| > |\tilde{q}_j|$  for at least one  $j$ , such that  $1 \leq j \leq d$ .

First, consider finite  $p \geq 1$ . By the properties of  $x^p$ , we have  $|q_j|^p \geq |\tilde{q}_j|^p \geq 0$  for all  $j$ . Furthermore,  $|q_j|^p > |\tilde{q}_j|^p$  for at least one  $j$ . This, in turn, implies

$$\sum_{j=1}^d |q_j|^p > \sum_{j=1}^d |\tilde{q}_j|^p \geq 0.$$

It follows that

$$\|q\|_p = \left( \sum_{j=1}^d |q_j|^p \right)^{1/p} > \left( \sum_{j=1}^d |\tilde{q}_j|^p \right)^{1/p} = \|\tilde{q}\|_p.$$

Consider the case  $p = \infty$  separately:

$$\|q\|_\infty = \lim_{p \rightarrow \infty} \left( \sum_{j=1}^d |q_j|^p \right)^{1/p} = \max_{1 \leq j \leq d} |q_j|.$$

We have  $\max_{1 \leq j \leq d} |q_j| > \max_{1 \leq j \leq d} |\tilde{q}_j|$ .

Therefore, for both the case where  $p \geq 1$  is finite and  $p = \infty$ ,  $\|\cdot\|_p$  is strictly one-sided monotone.  $\square$

Define a linearly-transformed  $p$ -norm,  $\|\cdot\|_{B,p}$ , to be

$$\|q\|_{B,p} = \|Bq\|_p,$$

where  $p \geq 1$  and  $B$  is a nonsingular  $d \times d$  matrix. Note that  $B$  must be nonsingular so that  $\|\cdot\|_{B,p}$  satisfies the properties of a norm such as definiteness and homogeneity. Such a norm is not strictly one-sided monotone in general. Figure 2.2(a) shows a simple example where a vector is rotated by  $-\pi/4$  and scaled by 3 in the  $q_2$ -axis before the Euclidean norm is taken; i.e.,

$$B = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} \cos(-\pi/4) & -\sin(-\pi/4) \\ \sin(-\pi/4) & \cos(-\pi/4) \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -3/\sqrt{2} & 3/\sqrt{2} \end{bmatrix}. \quad (2.3)$$

Let  $q = (2, 2)^T$  and  $\tilde{q} = (\sqrt{2}, 0)^T$ . We have  $q \succeq \tilde{q}$ , but,

$$\|Bq\|_2 = \|(2\sqrt{2}, 0)^T\|_2 = \sqrt{8} < \sqrt{10} = \|(1, -3)^T\|_2 = \|B\tilde{q}\|_2.$$

Consequently, this particular linearly-transformed  $p$ -norm is not strictly one-sided monotone. However, in this case an inverse transformation  $B^{-1}$  of the grid coordinates will result in a strictly one-sided monotone  $p$ -norm, while maintaining the grid's orthogonality. More generally, we conjecture that if the Hamiltonian is of the form  $H(q) = \tilde{H}(Bq)$ , where  $B$  is a rotation (which may be followed by scaling) and  $\tilde{H}$  satisfies Properties 1 to 4, a transformation of the grid coordinates by  $B^{-1}$  will result in a transformed  $H$  that also satisfies Properties 1 to 4, while maintaining the grid's orthogonality. More complex coordinate modifications might be possible but we have not yet adequately investigated conditions or procedures.

A scaled  $p$ -norm (Figure 2.2(b)) is a special case of a linearly-transformed  $p$ -norm. Such a norm scales the components of its argument before applying a  $p$ -norm,




FIG. 2.2. Contour plots of  $\|Bq\|_p$ . (a) is not strictly one-sided monotone:  $p = 2$  and  $B$  is defined by (2.3). (b) is strictly one-sided monotone:  $p = 1$  and  $B$  scales by 2 in the  $q_1$ -axis.




FIG. 2.3. Contour plots of  $G(q)$ . (a) mixed  $p$ -norm:  $G$  is defined by (2.4). (b) asymmetric norm-like function:  $G$  is defined by (2.5).

by restricting  $B$  to be a nonsingular diagonal matrix. It is simple to show that a scaled  $p$ -norm is strictly one-sided monotone, considering Proposition 2.6.

A mixed  $p$ -norm is a recursive composition of  $p$ -norms and it is strictly one-sided monotone. The following is an example (Figure 2.3(a)) of a mixed  $p$ -norm that takes the Euclidean norm of the first 2 components and then takes the Manhattan norm of the result and the last component:

$$\begin{aligned} \|q\| &= \|(\| (q_1, q_2) \|_2, q_3)\|_1 \\ &= \sqrt{(q_1)^2 + (q_2)^2} + |q_3|. \end{aligned} \quad (2.4)$$

where  $q = (q_1, q_2, q_3)$ . This particular norm was used as a  $G$  function in [1] for a simple 2-robot coordinated optimal control problem.

Finally, the one-sidedness of Property 4 allows  $G$  to be asymmetric, which is not permitted for a norm. An example of such an asymmetric norm-like function is shown

in Figure 2.3(b) and is given by

$$G(q) = \begin{cases} \|B_a q\|_\infty, & \text{if } q_1 \leq 0 \text{ and } q_2 \leq 0, \\ \|B_b q\|_1, & \text{if } q_1 \leq 0 \text{ and } q_2 > 0, \\ \|B_c q\|_2, & \text{if } q_1 > 0 \text{ and } q_2 \leq 0, \\ \|B_d q\|_2, & \text{if } q_1 > 0 \text{ and } q_2 > 0, \end{cases} \quad (2.5)$$

where

$$B_a = \begin{bmatrix} 1/2 & 0 \\ 0 & 1 \end{bmatrix} \quad B_b = \begin{bmatrix} 1/2 & 0 \\ 0 & 1/2 \end{bmatrix} \quad B_c = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad B_d = \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix}.$$

We solve the anisotropic problem characterized by (2.5) as well as an anelliptic wave propagation problem and a multi-robot optimal path planning problem in Section 5. Other examples of  $G$  functions which satisfy strict one-sided monotonicity are some polygonal norms such as axis aligned hexagonal or octagonal norms; however, we do not further investigate these options here.

**3. FMM and the Discretized Problem.** We define a discretized analogue of the Dirichlet problem (1.1). By describing the **Update** function in Algorithm 1, we also formalize the FMM algorithm. Finally, we examine important properties of the **Update** function and show that FMM solves the discretized problem.

We recall that the nodes in  $\underline{\Omega}$  lie on an axis-aligned orthogonal grid. Let  $\underline{x}_0 \in \underline{\Omega}$ . The neighborhood of  $\underline{x}_0$  is shown in Figure 3.1. Let  $\underline{x}_j^\pm$  be the neighbors of  $\underline{x}_0$  in the  $\pm e_j$  directions,  $e_j$  being the  $j$ th vector in the standard basis. The set of neighbors is

$$\mathcal{N}(\underline{x}_0) = \{\underline{x}_1^\pm, \underline{x}_2^\pm, \dots, \underline{x}_d^\pm\}$$

and the neighborhood vector is

$$N(\underline{x}_0) = (\underline{x}_0, \underline{x}_1^\pm, \underline{x}_2^\pm, \dots, \underline{x}_d^\pm).$$

Let  $h_j^\pm = \pm \|\underline{x}_0 - \underline{x}_j^\pm\|$  be signed distances to the neighbors in the  $\pm e_j$  directions. Let

$$\mathcal{S} = \{(s_1, s_2, \dots, s_d) \mid s_j \in \{-1, +1\}, 1 \leq j \leq d\},$$

such that  $s \in \mathcal{S}$  represents one of the  $2^d$  neighboring simplices of  $\underline{x}_0$ . Note that we abuse notation by using  $s_j \in \{-1, +1\}$  as a superscript indexing  $\underline{x}_j^\pm$  or  $h_j^\pm$ .

Let  $B(\Omega)$  be the set of bounded functions on domain  $\Omega$ . We define the numerical Hamiltonian  $\underline{H} : \Omega^{1+2d} \times B(\Omega) \times \mathbb{R} \rightarrow \mathbb{R}$  as follows:

$$\underline{H}(N, \phi, \mu) = \max_{s \in \mathcal{S}} [H(\underline{x}_0, \underline{D}^s(N, \phi, \mu))], \quad (3.1)$$

where  $H$  is as defined in Section 2 and

$$\underline{D}^s(N, \phi, \mu) = (\underline{D}_1^s(N, \phi, \mu), \underline{D}_2^s(N, \phi, \mu), \dots, \underline{D}_d^s(N, \phi, \mu))$$

is a first-order, upwind, finite-difference gradient approximation from the simplex represented by  $s$ ; that is,

$$\underline{D}_j^s(N, \phi, \mu) = \frac{\max(0, \mu - \phi(\underline{x}_j^{s_j}))}{-h_j^{s_j}}, \quad (3.2)$$




FIG. 3.1. Neighborhood of  $\underline{x}_0$  with  $d = 2$ .

for  $1 \leq j \leq d$ . Although  $\underline{H}$  is defined on domain  $\Omega^{1+2d} \times B(\Omega) \times \mathbb{R}$ , for FMM it will only be used on domain  $\underline{\Omega}^{1+2d} \times B(\underline{\Omega}) \times \mathbb{R}$ . The broader definition of domain is important for consistency [5]. The restriction of  $\Omega^{1+2d}$  to  $\underline{\Omega}^{1+2d}$  poses no problems to the definition of  $\underline{H}$ . Furthermore, to evaluate  $\underline{H}$ ,  $\phi$  need only be defined on  $\mathcal{N}$ , which is true of any function in  $B(\underline{\Omega})$ .

The discretized Dirichlet problem is to find a function  $\underline{u} : (\underline{\Omega} \cup \partial\underline{\Omega}) \rightarrow \mathbb{R}$ , such that

$$\underline{H}(N(\underline{x}), \underline{u}, \underline{u}(\underline{x})) = 0, \quad \underline{x} \in \underline{\Omega} \quad (3.3a)$$

$$\underline{u}(\underline{x}) = g(\underline{x}), \quad \underline{x} \in \partial\underline{\Omega}. \quad (3.3b)$$

DEFINITION 3.1. Let FMM be Algorithm 1 with the Update function defined as follows. A call to Update( $\underline{x}_0, \underline{u}$ ) returns the solution  $\mu = \tilde{\mu}$  to

$$\underline{H}(N(\underline{x}_0), \underline{u}, \mu) = 0. \quad (3.4)$$

In this way it determines a node's value  $\underline{u}(\underline{x}_0) \leftarrow \tilde{\mu}$  given the values of its neighbors,  $\underline{u}_j^\pm = \underline{u}(\underline{x}_j^\pm)$ . When we are varying only  $\mu$ , it will be convenient to write  $\underline{H}(\mu) = \underline{H}(N, \phi, \mu)$  and  $\underline{D}^s(\mu) = \underline{D}^s(N, \phi, \mu)$ . For the lemmas and theorems stated below we assume  $H$  satisfies Properties 1 to 4.

PROPOSITION 3.2. Let  $\underline{u} : (\underline{\Omega} \cup \partial\underline{\Omega}) \rightarrow \mathbb{R}$  be the grid function after FMM terminates. Then  $\underline{u}$  is the unique solution of (3.3).

This proposition states that the grid function  $\underline{u}$  that results from running FMM solves the discretized problem (3.3). It is proved in Section 3.4. For FMM, causality of the numerical scheme is essential, so we prove this property in Section 3.3.

A method for proving the convergence of  $\underline{u}$  to the solution of (1.1) as the grid spacing goes to zero is presented in [5]. It is shown there that the consistency, monotonicity and stability of the numerical scheme are sufficient for convergence. Also, uniqueness and monotonicity of the solution to (3.4) are useful for using numerical root finders to implement Update. We include proofs of uniqueness and monotonicity in Sections 3.1 and 3.2, respectively. More details regarding convergence, including the proofs of consistency and stability, are in Section 4.

**3.1. Unique Update.** Let the minimum value of all neighbours of  $\underline{x}_0$  be

$$\check{\underline{u}} = \min_{x \in \mathcal{N}(\underline{x}_0)} (\underline{u}(x)). \quad (3.5)$$

We show there is a unique solution  $\mu = \tilde{\mu}$  to (3.4), such that  $\tilde{\mu} > \check{\underline{u}}$ . First, we prove two useful lemmas.

LEMMA 3.3.  $\underline{H}(\mu)$  is strictly increasing on  $\mu \geq \check{\underline{u}}$ .

*Proof.* Let  $\mu_a > \mu_b \geq \check{\underline{u}}$ . Let  $s \in \mathcal{S}$  and  $1 \leq j \leq d$ . If  $\mu_a > \underline{u}_j^{s_j}$  then  $\underline{D}_j^s(\mu_a)\underline{D}_j^s(\mu_b) \geq 0$  and  $|\underline{D}_j^s(\mu_a)| > |\underline{D}_j^s(\mu_b)|$ . On the other hand, if  $\mu_a \leq \underline{u}_j^{s_j}$  then  $\underline{D}_j^s(\mu_a) = \underline{D}_j^s(\mu_b) = 0$ . Also, there exists at least one  $s \in \mathcal{S}$  and  $1 \leq j \leq d$  such that  $\underline{D}_j^s(\mu_a) \neq 0$ , since  $\mu_a > \check{\underline{u}}$ . For such  $s$ ,  $H(\underline{D}^s(\mu_a)) > H(\underline{D}^s(\mu_b))$ , by strict one-sided monotonicity (Property 4). For all other  $s$ ,  $H(\underline{D}^s(\mu_a)) = H(\underline{D}^s(\mu_b)) = H(0)$ . Therefore, by (3.1)  $\underline{H}(\mu_a) > \underline{H}(\mu_b)$ , so  $\underline{H}(\mu)$  is strictly increasing on  $\mu \geq \check{\underline{u}}$ .  $\square$

LEMMA 3.4. The numerical Hamiltonian  $\underline{H}(\mu)$  satisfies the following.

- (a)  $\underline{H}(\mu) = H(0) < 0$  for  $\mu \leq \check{\underline{u}}$ .
- (b)  $\underline{H}(\mu) \rightarrow \infty$  as  $\mu \rightarrow \infty$ .
- (c)  $\underline{H}(\mu)$  is nondecreasing on all  $\mu$ .

*Proof.* If  $\mu \leq \check{\underline{u}}$  then by (3.2) and (3.5), we have  $\underline{D}_j^s(\mu) = 0$  for all  $s \in \mathcal{S}$ ,  $1 \leq j \leq d$ . By the strict compatibility of  $H$ ,  $H(\underline{D}^s(v_j)) = H(0) < 0$ , for all  $s$ . By (3.1), we have  $\underline{H}(\mu) = H(0) < 0$ , for  $\mu \leq \check{\underline{u}}$ , proving (a).

Let  $s \in \mathcal{S}$  and  $1 \leq j \leq d$ . As  $\mu \rightarrow \infty$ , we have  $\underline{D}_j^s(\mu) \rightarrow \infty$  and  $\|\underline{D}^s(\mu)\| \rightarrow \infty$  for all  $s \in \mathcal{S}$ ,  $1 \leq j \leq d$ . By the coercivity of  $H$ , as  $\mu \rightarrow \infty$ , we have  $H(\underline{D}^s(\mu)) \rightarrow \infty$  for all  $s \in \mathcal{S}$ . By (3.1), we have  $\underline{H}(\mu) \rightarrow \infty$  as  $\mu \rightarrow \infty$ , proving (b).

Because  $\underline{H}(\mu)$  is constant on  $\mu \leq \check{\underline{u}}$  and by Lemma 3.3 increasing on  $\mu \geq \check{\underline{u}}$ ,  $\underline{H}(\mu)$  is nondecreasing on all  $\mu$ , proving (c).  $\square$

THEOREM 3.5. There exists a unique solution  $\mu = \tilde{\mu}$  to  $\underline{H}(\mu) = 0$  such that  $\tilde{\mu} > \check{\underline{u}}$ .

*Proof.* Each  $\underline{D}_j^s(\mu)$  is continuous on  $\mu$ . Furthermore, by the continuity of  $H$ ,  $H(\underline{D}^s(\mu))$  is continuous on  $\mu$ , for all  $s$ . Since max is continuous,  $\underline{H}(\mu)$  is continuous. By Lemma 3.4(a/b),  $\underline{H}(\mu) < 0$  for  $\mu \leq \check{\underline{u}}$  and  $\underline{H}(\mu) \rightarrow \infty$  as  $\mu \rightarrow \infty$ . Therefore, by the Intermediate Value Theorem there exists a solution  $\mu = \tilde{\mu}$  to  $\underline{H}(\mu) = 0$ , such that  $\check{\underline{u}} < \tilde{\mu} < \infty$ . Moreover, since  $\underline{H}$  is strictly increasing on  $\mu \geq \check{\underline{u}}$  by Lemma 3.3, the solution is unique.  $\square$

REMARK 1. We note that strict one-sided monotonicity (Property 4) of  $H$  is used to prove Lemma 3.3, and Lemma 3.3 is then used to show that the solution to  $\underline{H}(\mu) = 0$  is unique. We might consider whether or not one-sided monotonicity (Property 5) of  $H$  is sufficient for a unique solution. However, Property 5 would not be sufficient to prove Lemma 3.3 and we would find that  $\underline{H}(\mu)$  is only nondecreasing on  $\mu \geq \check{\underline{u}}$ . A solution to  $\underline{H}(\mu) = 0$  would still be guaranteed but not unique in this case. Analogously, for differentiable  $H$ , Osher's criterion on  $H$  implies a solution that may not be unique unless  $H$  satisfies some additional property, such as one-sided homogeneity (Property 6) or convexity.

**3.2. Monotonicity.** We show that  $\underline{H}$  and the Update function are monotone in the neighbor's values. Monotonicity of  $\underline{H}$  requires that if none of the neighbor's values decreases, the numerical Hamiltonian  $\underline{H}$  should not increase. Additionally, monotonicity of the Update function requires that if none of the neighbor's values decreases, the solution to (3.4) should not decrease. Monotonicity is useful both for showing that FMM finds a unique solution and for proving convergence. We note

that monotonicity does not require strict one-sided monotonicity of  $H$ , but rather one-sided monotonicity of  $H$  is sufficient.

**THEOREM 3.6.** *Let  $\underline{v}$  and  $\underline{u}$  be grid functions. Let  $v_j^\pm \geq u_j^\pm$  for  $1 \leq j \leq d$ . Then for  $\mu \in \mathbb{R}$ , we have  $\underline{H}(N, \underline{v}, \mu) \leq \underline{H}(N, \underline{u}, \mu)$ . Furthermore, if  $\mu = \mu_v$  is the unique solution to  $\underline{H}(N, \underline{v}, \mu) = 0$  and  $\mu = \mu_u$  is the unique solution to  $\underline{H}(N, \underline{u}, \mu) = 0$ , then  $\mu_v \geq \mu_u$ .*

*Proof.* Let  $\mu \in \mathbb{R}$ . We have  $\underline{D}^s(N, \underline{u}, \mu) \supseteq \underline{D}^s(N, \underline{v}, \mu)$ , for all  $s \in \mathcal{S}$ . Also, by Proposition 2.3,  $H$  satisfies one-sided monotonicity (Property 5). Thus,

$$H(\underline{D}^s(N, \underline{u}, \mu)) \geq H(\underline{D}^s(N, \underline{v}, \mu)) = 0,$$

for all  $s \in \mathcal{S}$ . Consequently,  $\underline{H}(N, \underline{u}, \mu) \geq \underline{H}(N, \underline{v}, \mu)$ , proving the first claim.

To prove the second claim, we let  $\mu_v$  and  $\mu_u$  be as defined above. We note that  $\underline{H}(N, \underline{u}, \mu_u) = 0 \geq \underline{H}(N, \underline{v}, \mu_u)$ . By Lemma 3.4(c),  $\underline{H}(N, \underline{v}, \mu)$  is nondecreasing on all  $\mu$ , so in order that  $\underline{H}(N, \underline{v}, \mu_v) = 0$ , it must be that  $\mu_v \geq \mu_u$ .  $\square$

**3.3. Causality.** We note that (3.3) defines a very large system of nonlinear equations, one equation for each node  $\underline{x} \in \underline{\Omega}$ . FMM can be used to solve this system very efficiently, if the solution  $\mu = \tilde{\mu}$  to (3.4) is dependent only on neighbors with smaller values. This property represents a causal relationship between node values. There is an information flow from nodes with smaller values to those with larger values. The causal relationship is meant to mimic that of the PDE (1.1). The solution  $u$  of (1.1) is completely defined at  $x$  using only values of  $u$  from states that are backwards along the characteristic line that passes through  $x$ .

FMM exploits the causal property of  $\underline{H}$  by computing  $\underline{u}(\underline{x})$  in increasing order in a single pass through the nodes. This causal property has been discussed as a requirement for Dijkstra-like single-pass methods in several works [30, 23, 25, 19, 28]. The following theorem states that  $\underline{H}$  and the **Update** function are causal. The **Update** function is considered causal if any change to the value of a neighbouring node, such that the both the new and old values are no smaller than the solution  $\mu = \tilde{\mu}$  to  $\underline{H}(N, \underline{u}, \mu) = 0$ , has no effect on the solution.

**THEOREM 3.7.** *Let  $\underline{v}$  and  $\underline{u}$  be grid functions. Let*

$$\tilde{\mathcal{N}}(\underline{x}_0) = \{\underline{x} \in \mathcal{N}(\underline{x}_0) \mid \underline{v}(\underline{x}) \neq \underline{u}(\underline{x})\}.$$

*Let*

$$\tilde{w} = \begin{cases} \min_{\underline{x} \in \tilde{\mathcal{N}}(\underline{x}_0)} \min(\underline{v}(\underline{x}), \underline{u}(\underline{x})), & \text{if } \tilde{\mathcal{N}}(\underline{x}_0) \neq \emptyset, \\ +\infty, & \text{otherwise.} \end{cases}$$

*Then  $\underline{H}(N, \underline{v}, \mu) = \underline{H}(N, \underline{u}, \mu)$ , for  $\mu \leq \tilde{w}$ .*

*Furthermore, let  $\mu = \tilde{\mu}_u$  be the unique solution to  $\underline{H}(N, \underline{u}, \mu) = 0$ , and  $\mu = \tilde{\mu}_v$  be the unique solution to  $\underline{H}(N, \underline{v}, \mu) = 0$ . If  $\tilde{\mu}_u \leq \tilde{w}$  or  $\tilde{\mu}_v \leq \tilde{w}$  then  $\tilde{\mu}_u = \tilde{\mu}_v$ .*

*Proof.* Let  $\mu \leq \tilde{w}$ . By (3.2) and the definition of  $\tilde{w}$ , we have  $D_j^s(N, \underline{v}, \mu) = D_j^s(N, \underline{u}, \mu)$ , for all  $s \in \mathcal{S}$ ,  $1 \leq j \leq d$ . This implies that  $\underline{H}(N, \underline{v}, \mu) = \underline{H}(N, \underline{u}, \mu)$ , proving the first claim.

For the second claim, let  $\tilde{\mu}_u$  and  $\tilde{\mu}_v$  be as defined above. Let  $\tilde{\mu}_u \leq \tilde{w}$ . Then  $\underline{H}(N, \underline{v}, \tilde{\mu}_u) = \underline{H}(N, \underline{u}, \tilde{\mu}_u) = 0$ , so  $\mu = \tilde{\mu}_u$  is a solution to  $\underline{H}(N, \underline{v}, \mu) = 0$ . By Theorem 3.5, this solution is unique. By a symmetric argument, if  $\tilde{\mu}_v \leq \tilde{w}$  then  $\mu = \tilde{\mu}_v$  is the unique solution to  $\underline{H}(N, \underline{u}, \mu) = 0$ .  $\square$

**3.4. Solution.** We show that FMM finds a unique solution to (3.3). The proof method in this section is similar to those for isotropic FMM in [31, 22]. The causality of the `Update` function proved in Section 3.3 is key to the proof. We begin by proving the following lemmas: Lemma 3.8 states that Algorithm 1 terminates and Lemma 3.9 states that the grid function does not increase as Algorithm 1 progresses.

LEMMA 3.8. *Let  $|\underline{\Omega} \cup \partial \underline{\Omega}|$  be finite. If `ExtractMin` and `Update` always terminate then Algorithm 1 terminates with at most  $2d|\underline{\Omega} \cup \partial \underline{\Omega}|$  calls to `Update`.*

*Proof.* Since  $\mathcal{Q}$  is initialized to  $\underline{\Omega} \cup \partial \underline{\Omega}$  in Line 3 of Algorithm 1, it has a finite number of elements. For every iteration of the **while** loop one element is removed from  $\mathcal{Q}$  and at most  $2d$  neighbors are updated. The **while** loop terminates when  $\mathcal{Q} = \emptyset$ . Therefore, there are at most  $|\underline{\Omega} \cup \partial \underline{\Omega}|$  iterations of the **while** loop and at most  $2d|\underline{\Omega} \cup \partial \underline{\Omega}|$  calls to `Update`.  $\square$

As Algorithm 1 is run the grid function  $\underline{u}$  evolves. Accordingly, it is useful to talk about a sequence of grid functions  $\underline{u}^k$ . Let  $\underline{u}^0$  be the state of  $\underline{u}$  just after initialization on Line 3 in Algorithm 1. Let  $\underline{u}^k$  for  $k \geq 1$  be the state of the grid function after the  $k$ th call to `Update`. For two grid functions  $\underline{u}$  and  $\tilde{u}$ , we say that  $\underline{u} \leq \tilde{u}$  if and only if  $\underline{u}(\underline{x}) \leq \tilde{u}(\underline{x})$  for all  $\underline{x} \in (\underline{\Omega} \cup \partial \underline{\Omega})$ .

LEMMA 3.9. *The sequence of grid functions  $\underline{u}^k$  is nonincreasing on  $k$ .*

*Proof.* Consider a node  $\underline{x} \in \underline{\Omega}$  being updated using

$$\underline{u}^{k'+1}(\underline{x}) \leftarrow \text{Update}(\underline{x}, \underline{u}^{k'}).$$

Assume that the subsequence  $\underline{u}^l$  is nonincreasing on  $0 \leq l \leq k'$ . Either  $\underline{x}$  has been updated before or it has not and  $\underline{u}^{k'}(\underline{x}) = \infty$ . In the latter case, certainly  $\underline{u}^{k'+1}(\underline{x}) \leq \underline{u}^{k'}(\underline{x}) = \infty$ . In the former case, let the previous update of  $\underline{x}$  be

$$\underline{u}^{k''+1}(\underline{x}) \leftarrow \text{Update}(\underline{x}, \underline{u}^{k''}),$$

where  $k'' < k'$ . By the assumption,  $\underline{u}^{k'} \leq \underline{u}^{k''}$ . By Theorem 3.6,

$$\underline{u}^{k'+1}(\underline{x}) \leq \underline{u}^{k''+1}(\underline{x}) = \underline{u}^{k'}(\underline{x}).$$

Consequently,  $\underline{u}^{k'+1} \leq \underline{u}^{k'}$ . We also have  $\underline{u}^1 \leq \underline{u}^0$  since the first update to any node  $\underline{x} \in \underline{\Omega}$  cannot be such that  $\underline{u}^1(\underline{x}) > \underline{u}^0(\underline{x}) = \infty$ . Therefore, by induction on  $k$ ,  $\underline{u}^k$  is a nonincreasing sequence.  $\square$

PROPOSITION 3.10. *Let  $\underline{u} : (\underline{\Omega} \cup \partial \underline{\Omega}) \rightarrow \mathbb{R}$  be the grid function after FMM terminates. Then  $\underline{u}$  is the unique solution of (3.3).*

If a node  $\underline{x} \notin \mathcal{Q}$  we say that it is *known*, meaning that its value can no longer be changed by Algorithm 1. If a node  $\underline{x} \in \mathcal{Q}$  we say that it is *estimated*, meaning that its value can still be changed. When a node  $\underline{x}$  is extracted from  $\mathcal{Q}$  we say that it *becomes known*. Let  $\mathcal{Q}^l$  be a sequence of node sets such that  $\mathcal{Q}^0 = \underline{\Omega} \cup \partial \underline{\Omega}$  and  $\mathcal{Q}^l$  for  $l \geq 1$  is the state of  $\mathcal{Q}$  after the  $l$ th call to `ExtractMin`.

We prove the theorem in two stages. First, we show that when a node  $\underline{x}$  *becomes known*, (3.3) is satisfied for that node. Then, we show that after  $\underline{x}$  *becomes known* only neighbors' values not less than  $\underline{u}(\underline{x})$  may change and the changed neighbors' values will not be less than  $\underline{u}(\underline{x})$ . Using Theorem 3.7, (3.3) remains satisfied for  $\underline{x}$  that is *known*.

*Proof.* Consider a node  $\underline{x} \in \underline{\Omega}$  that has just become known, such that  $\mathcal{Q}^{l'+1} = \mathcal{Q}^l \setminus \{\underline{x}\}$ . Let  $\underline{u}^{k'}$  be the grid function when  $\underline{x}$  becomes known. Since  $\underline{u}^{k'}(\underline{x})$  can no longer change, we say  $\underline{u}(\underline{x}) = \underline{u}^{k'}(\underline{x})$ . Let the previous update of  $\underline{x}$  be

$$\underline{u}^{k''+1}(\underline{x}) \leftarrow \text{Update}(\underline{x}, \underline{u}^{k''}),$$



where  $k'' + 1 \leq k'$ . Note that  $\underline{u}(\underline{x}) = \underline{u}^{k''+1}(\underline{x})$  and

$$\underline{H}(N(\underline{x}), \underline{u}^{k''+1}, \underline{u}^{k''+1}(\underline{x})) = 0.$$

Let  $\mathcal{Q}^{l''}$  be the state of  $\mathcal{Q}$  at the time of this previous update. Since the previous update of  $\underline{x}$  occurred just after the last time a neighbor  $\tilde{x} \in \mathcal{N}(\underline{x})$  became known, we have  $\mathcal{Q}^{l'} \cup \mathcal{N}(\underline{x}) = \mathcal{Q}^{l''} \cup \mathcal{N}(\underline{x})$ . But any neighbor  $\tilde{x} \in \mathcal{N}(\underline{x})$  for which  $\underline{u}^{k'}(\tilde{x}) \neq \underline{u}^{k''+1}(\tilde{x})$  must be in  $\mathcal{Q}^{l'} \cup \mathcal{N}(\underline{x})$ , since only *estimated* nodes are updated. Also, since **ExtractMin** removes the *estimated* node with minimum value,  $\underline{u}^{k'}(\tilde{x}) \geq \underline{u}(\underline{x})$ , for all  $\tilde{x} \in \mathcal{Q}^{l'} \cup \mathcal{N}(\underline{x})$ . Furthermore,  $\underline{u}^{k''+1}(\tilde{x}) \geq \underline{u}(\underline{x})$ , for all  $\tilde{x} \in \mathcal{Q}^{l'} \cup \mathcal{N}(\underline{x})$ , since  $\underline{u}^{k''+1} \geq \underline{u}^{k'}$  by Lemma 3.9. By Theorem 3.7,

$$\underline{H}(N(\underline{x}), \underline{u}^{k'}, \underline{u}(\underline{x})) = \underline{H}(N(\underline{x}), \underline{u}^{k''+1}, \underline{u}^{k''+1}(\underline{x})) = 0.$$

Now consider the  $i$ th neighbor  $\underline{x}^i \in \mathcal{N}(\underline{x}) \cap \mathcal{Q}^{l'+1}$  updated after  $\underline{x}$  became known but before any other *estimated* nodes become known with

$$\underline{u}^{k'+i}(\underline{x}^i) \leftarrow \text{Update}(\underline{x}^i, \underline{u}^{k'+i-1}).$$

Let the previous update of  $\underline{x}^i$  be

$$\underline{u}^{k''' + 1}(\underline{x}^i) \leftarrow \text{Update}(\underline{x}^i, \underline{u}^{k'''}), \quad (3.6)$$

where  $k''' + 1 \leq k' \leq k' + i - 1$ . Assume  $\underline{u}^{k'+i-1}(y) \geq \underline{u}(\underline{x})$ , for all  $y \in \mathcal{Q}^{l'}$ . We also have  $\underline{u}^{k''' + 1}(y) \geq \underline{u}(\underline{x})$ , for all  $y \in \mathcal{Q}^{l'}$ , since  $\underline{u}^{k''' + 1} \geq \underline{u}^{k'+i-1}$  by Lemma 3.9.

Let  $\tilde{x} \in \mathcal{N}(\underline{x}^i)$  be such that  $\underline{u}^{k'+i-1}(\tilde{x}) \neq \underline{u}^{k''' + 1}(\tilde{x})$ . Such  $\tilde{x}$  could not have become known before  $\underline{x}$  became known, because the previous update (3.6) of  $\underline{x}^i$  would follow and that would imply  $\underline{u}^{k'+i-1}(\tilde{x}) = \underline{u}^{k''' + 1}(\tilde{x})$ , a contradiction. Consequently, if  $\tilde{x} \in \mathcal{N}(\underline{x}^i)$  and  $\underline{u}^{k'+i-1}(\tilde{x}) \neq \underline{u}^{k''' + 1}(\tilde{x})$ , then  $\tilde{x} \in \mathcal{Q}^{l'}$  and  $\underline{u}^{k''' + 1}(\tilde{x}) > \underline{u}^{k'+i-1}(\tilde{x}) \geq \underline{u}(\underline{x})$ . By Theorem 3.7, if  $\underline{u}^{k'+i}(\underline{x}^i) \neq \underline{u}^{k''' + 1}(\underline{x}^i)$ , then  $\underline{u}^{k'+i}(\underline{x}^i) \geq \underline{u}(\underline{x})$ . On the other hand, we have  $\underline{u}^{k'+i}(\underline{x}^i) = \underline{u}^{k'+i-1}(\underline{x}^i)$  and by the assumption  $\underline{u}^{k'+i-1}(\underline{x}^i) \geq \underline{u}(\underline{x})$ . So, for all  $y \in \mathcal{Q}^{l'}$ ,  $\underline{u}^{k'+i}(y) \geq \underline{u}(\underline{x})$ .

We also know that  $\underline{u}^{k'}(y) \geq \underline{u}(\underline{x})$ , for all  $y \in \mathcal{Q}^{l'}$ , since **ExtractMin** removes the *estimated* node with minimum value. By induction on  $i$ ,  $\underline{u}^{k'+i}(y) \geq \underline{u}(\underline{x})$ , for all  $y \in \mathcal{Q}^{l'}$ , for all  $i$  such that  $1 \leq i \leq |\mathcal{N}(\underline{x}) \cap \mathcal{Q}^{l'+1}|$ .

By induction on the sequence of nodes to become known after  $\underline{x}$  becomes known we find that any such node  $\underline{y}$  must be such that  $\underline{u}(\underline{y}) \geq \underline{u}(\underline{x})$ . Note that only nodes  $\underline{y}$  that become known after  $\underline{x}$  becomes known may have values such that  $\underline{u}(\underline{y}) \neq \underline{u}^{k'}(\underline{y})$ . Therefore, by Theorem 3.7,

$$\underline{H}(N(\underline{x}), \underline{u}, \underline{u}(\underline{x})) = \underline{H}(N(\underline{x}), \underline{u}^{k'}, \underline{u}(\underline{x})) = 0.$$

Since, by Lemma 3.8, every node  $\underline{x} \in \underline{\Omega} \cup \partial \underline{\Omega}$  becomes known in the course of Algorithm 1,  $\underline{u}$  is a solution of (3.3). By Theorem 3.5, this solution is unique.  $\square$

**4. Convergence.** For convergence of FMM-computed  $\underline{u}$  to the viscosity solution  $u$  of (1.1), an update scheme must be consistent, monotone, and stable [5]. Monotonicity is proven in Section (3.2). We closely follow the technique described in [5] to prove convergence in this section.

**4.1. Consistency.** We show that the numerical Hamiltonian  $\underline{H}$  is consistent with (2.2). Let  $C_b^\infty(\Omega)$  be the set of smooth, bounded functions on domain  $\Omega$ .

**THEOREM 4.1.** *Let  $\phi \in C_b^\infty(\Omega)$ . Let  $x \in \Omega$ . Let  $H$  be continuous in the first argument and satisfy Properties 1 to 4. Then*

$$\lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0 \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} \underline{H}(N, \phi + \xi, \phi(\underline{x}_0) + \xi) = H(x, D\phi(x)). \quad (4.1)$$

*Proof.* Let  $\phi$ ,  $x$ , and  $H$  be as defined above. Let

$$D\phi(x) = (\partial_1\phi(x), \partial_2\phi(x), \dots, \partial_d\phi(x)).$$

Let  $s \in \mathcal{S}$  and  $1 \leq j \leq d$ . We have by (3.2) and the smoothness of  $\phi$

$$\begin{aligned} \lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0}} D_j^s(N, \phi + \xi, \phi(\underline{x}_0) + \xi) &= \lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0}} \frac{\max(0, \phi(\underline{x}_0) + \xi - \phi(\underline{x}_j^{s_j}) - \xi)}{-h_j^{s_j}} \\ &= \lim_{\substack{\underline{x}_0 \rightarrow x \\ h_j^{s_j} \rightarrow 0}} \frac{\min(0, \phi(\underline{x}_0 + h_j^{s_j} e_j) - \phi(\underline{x}_0))}{h_j^{s_j}} \\ &= \begin{cases} \partial_j\phi(x), & \text{if } s_j\partial_j\phi(x) \leq 0, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Define  $\tilde{s}$  as

$$\tilde{s}_j = \begin{cases} +1, & \text{if } \partial_j\phi(x) \leq 0, \\ -1, & \text{otherwise,} \end{cases} \quad (4.2)$$

for  $1 \leq j \leq d$ . We have

$$\lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} D^{\tilde{s}}(N, \phi + \xi, \phi(\underline{x}_0) + \xi) = D\phi(x). \quad (4.3)$$

By the continuity and strict one-sided monotonicity of  $H$ ,

$$\lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} H(\underline{x}_0, D^{\tilde{s}}(N, \phi + \xi, \phi(\underline{x}_0) + \xi)) \geq \lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} H(\underline{x}_0, D^s(N, \phi + \xi, \phi(\underline{x}_0) + \xi)),$$

for all  $s$ . Therefore, by the continuity of  $\max$  and  $H$

$$\begin{aligned} &\lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} \underline{H}(N, \phi + \xi, \phi(\underline{x}_0) + \xi) \\ &= \lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} \left[ \max_{s \in \mathcal{S}} H(\underline{x}_0, D^s(N, \phi + \xi, \phi(\underline{x}_0) + \xi)) \right] \\ &= \lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} [H(\underline{x}_0, D^{\tilde{s}}(N, \phi + \xi, \phi(\underline{x}_0) + \xi))] \\ &= H(x, D\phi(x)). \end{aligned}$$

□

**4.2. Stability.** We show that for an orthogonal discretization  $\underline{\Omega}$ , the solution  $\underline{u}^\Omega$  to the discretized problem is bounded. We begin by showing with a lemma that the magnitude of the slope in  $\underline{u}^\Omega$  as measured between two neighbors is bounded.

LEMMA 4.2. *Let  $\underline{x}_0 \in \underline{\Omega}$  and  $\underline{H}(N(\underline{x}_0), \underline{u}, \mu) = 0$ . There exists  $\hat{K} < \infty$  such that*

$$|D_j^s(N(\underline{x}_0), \underline{u}, \mu)| \leq \hat{K},$$

for all  $s \in \mathcal{S}$ ,  $1 \leq j \leq d$ , and  $\underline{x}_0 \in \underline{\Omega}$ , such that  $\underline{H}(N(\underline{x}_0), \underline{u}, \mu) = 0$

*Proof.* Assume there exists some  $s \in \mathcal{S}$ ,  $1 \leq j \leq d$ , and  $\underline{x}_0 \in \underline{\Omega}$  with  $\underline{H}(N(\underline{x}_0), \underline{u}, \mu) = 0$ , such that  $|D_j^s(N(\underline{x}_0), \underline{u}, \mu)| = \infty$ . But by (3.1) and strict one-sided monotonicity and coercivity of  $H$ ,

$$\begin{aligned} \underline{H}(N(\underline{x}_0), \underline{u}, \mu) &\geq H(\underline{x}_0, D^s(N(\underline{x}_0), \underline{u}, \mu)) \\ &\geq H(\underline{x}_0, -|D_j^s(N(\underline{x}_0), \underline{u}, \mu)|s_j e_j) = \infty, \end{aligned}$$

contradicting the fact that  $\underline{H}(N(\underline{x}_0), \underline{u}, \mu) = 0$ . □

Let  $\underline{X} = (\underline{x}_1, \underline{x}_2, \dots, \underline{x}_k)$  be a sequence of neighboring nodes, such that  $\underline{x}_l \in \underline{\Omega} \cup \partial \underline{\Omega}$  for  $1 \leq l \leq k$  and  $\underline{x}_l \in \mathcal{N}(\underline{x}_{l-1})$  for  $2 \leq l \leq k$ . Define the grid path length of  $\underline{X}$  by

$$\omega(\underline{X}) = \sum_{l=2}^k \|\underline{x}_l - \underline{x}_{l-1}\|_2.$$

Define the minimum grid path length between  $\underline{x}$  and  $\underline{x}'$  to be

$$\tilde{\omega}(\underline{x}, \underline{x}') = \min\{\omega(\underline{X}) \mid \underline{x}_1 = \underline{x} \text{ and } \underline{x}_k = \underline{x}'\}.$$

Finally, define the minimum node-to-boundary grid path length of  $\underline{x}$  as

$$\tilde{\omega}(\underline{x}) = \min_{\underline{x}' \in \partial \underline{\Omega}} \tilde{\omega}(\underline{x}, \underline{x}').$$

Let  $\underline{u}^\Omega : \underline{\Omega} \cup \partial \underline{\Omega} \rightarrow \mathbb{R}$  be the solution to the discretized problem computed by FMM on the grid  $\underline{\Omega}$ . Let  $u^\Omega : \Omega \rightarrow \mathbb{R}$  extend  $\underline{u}^\Omega$  to the continuous domain using some non-expansive interpolator, such as nearest-neighbor or linear interpolation, for values between nodes. We show that for a reasonable discretization, such that the minimum distance along grid lines from any node to the boundary is bounded, the solution  $u^\Omega$  is bounded.

THEOREM 4.3. *Let  $\underline{\Omega}$  and  $\partial \underline{\Omega}$  be an orthogonal discretization such that for all  $\underline{x} \in \underline{\Omega}$ ,  $\tilde{\omega}(\underline{x}) \leq \hat{W}$ , for some constant  $\hat{W}$ . Then  $\min_{\underline{x}' \in \partial \underline{\Omega}} g(\underline{x}') \leq u^\Omega \leq \hat{U}$ , for some constant  $\hat{U}$ .*

*Proof.* Let  $\underline{x} \in \underline{\Omega}$ . Let  $\underline{x}' \in \partial \underline{\Omega}$  and  $\underline{X} = (\underline{x}_1, \underline{x}_2, \dots, \underline{x}_k)$  be such that  $\underline{x}_1 = \underline{x}$ ,  $\underline{x}_k = \underline{x}'$ , and  $\omega(\underline{X}) = \tilde{\omega}(\underline{x})$ . Obtain a modified discretization defined by

$$\begin{aligned} \tilde{\Omega} &= \{\underline{x} \mid \underline{x} \in \underline{\Omega} \text{ and } \underline{x} = \underline{x}_j, \text{ for some } j \text{ such that } 1 \leq j \leq k\} \text{ and} \\ \partial \tilde{\Omega} &= (\underline{\Omega} \cup \partial \underline{\Omega}) \setminus \tilde{\Omega}. \end{aligned}$$

Define the boundary condition  $\tilde{g} : \partial \tilde{\Omega} \rightarrow \mathbb{R}$  for the modified discretization

$$\tilde{g}(\underline{x}) = \begin{cases} g(\underline{x}), & \underline{x} \in \partial \underline{\Omega}, \\ \infty, & \text{otherwise.} \end{cases}$$

Run FMM on the modified grid and let  $\underline{u}^{\hat{\Omega}}(\underline{x})$  be the value computed at  $\underline{x}$ . Note that  $\underline{u}^{\hat{\Omega}}(\underline{x}) \leq \hat{W}\hat{K} + g(\underline{x}')$ , where  $\hat{K}$  is given by Lemma 4.2 and  $g$  is the original boundary condition. Also note, by the monotonicity of  $\underline{H}$  (Theorem 3.6), we have  $\underline{u}^{\hat{\Omega}}(\underline{x}) \leq \underline{u}^{\Omega}(\underline{x})$ , where  $\underline{u}^{\Omega}(\underline{x})$  is the value computed at  $\underline{x}$  by FMM on the original grid defined by  $\Omega$  and  $\partial\Omega$ . Therefore,  $\underline{u}^{\hat{\Omega}} \leq \hat{W}\hat{K} + \max_{\underline{x}' \in \partial\Omega} g(\underline{x}')$ . Since  $u^{\Omega}$  is defined by a non-expansive interpolation of the node values of  $\underline{u}^{\hat{\Omega}}$ ,

$$u^{\Omega} \leq \hat{W}\hat{K} + \max_{\underline{x}' \in \partial\Omega} g(\underline{x}').$$

For the lower bound, by Theorem 3.5, any solution  $\mu = \tilde{\mu}$  to  $\underline{H}(N(\underline{x}_0), \underline{u}, \mu) = 0$  must be such that  $\tilde{\mu} > \underline{u}$ , where  $\underline{u}$  is the minimum neighbor value (3.5). By induction on the nodes of  $\underline{\Omega}$ , we find that for any node  $\underline{x} \in \underline{\Omega} \cup \partial\underline{\Omega}$ ,

$$\underline{u}^{\Omega}(\underline{x}) \geq \min_{\underline{x}' \in \partial\underline{\Omega}} g(\underline{x}').$$

Because of the non-expansive interpolation to form  $u^{\Omega}$ , it follows that

$$u^{\Omega} \geq \min_{\underline{x}' \in \partial\underline{\Omega}} g(\underline{x}').$$

□

**4.3. Convergence.** Let  $\hat{h}(\underline{\Omega})$  be the maximum grid spacing between any two neighbors in  $\underline{\Omega}$ , i.e.,

$$\hat{h}(\underline{\Omega}) = \max_{\underline{x} \in \underline{\Omega}, \underline{x}' \in \mathcal{N}(\underline{x})} \|\underline{x} - \underline{x}'\|.$$

We aim to show that  $u^{\Omega}$  converges to the viscosity solution of (1.1) as  $\hat{h}(\underline{\Omega}) \rightarrow 0$ . An upper (respectively, lower) semi-continuous function  $u$  is a viscosity subsolution (respectively, supersolution) of (1.1) if for all  $\phi \in C_b^{\infty}(\Omega)$  such that  $u - \phi$  has a local maximum (respectively, minimum) at  $x$  we have

$$H(x, D\phi(x)) \leq 0 \quad (\text{respectively, } H(x, D\phi(x)) \geq 0). \quad (4.4)$$

A continuous function  $u$  is a viscosity solution if it is both a viscosity subsolution and a viscosity supersolution.

Assume any orthogonal discretization of  $\Omega$  and  $\partial\Omega$  is reasonable, as required by Theorem 4.3. We then have bounded  $u^{\Omega}$  and may define  $\hat{u}, \check{u} \in B(\Omega)$  by

$$\hat{u}(x) = \limsup_{\substack{\underline{x}' \rightarrow x \\ \hat{h}(\underline{\Omega}) \rightarrow 0}} u^{\Omega}(\underline{x}') \quad \text{and} \quad \check{u}(x) = \liminf_{\substack{\underline{x}' \rightarrow x \\ \hat{h}(\underline{\Omega}) \rightarrow 0}} u^{\Omega}(\underline{x}'). \quad (4.5)$$

Note that  $\hat{u}$  is upper semi-continuous and  $\check{u}$  is lower semi-continuous. The following proofs closely follow the exposition of [5].

**THEOREM 4.4.**  *$\hat{u}$  is a viscosity subsolution of (1.1) and  $\check{u}$  is a viscosity supersolution of (1.1).*

We only prove that  $\hat{u}$  is a viscosity subsolution of (1.1), as the second part of the theorem can be proved symmetrically.

*Proof.* Let  $\hat{x} \in \Omega$  be a local maximum of  $\hat{u} - \phi$  for some  $\phi \in C_b^{\infty}(\Omega)$ . Without loss of generality, assume  $\hat{x}$  is a strict local maximum [11, p. 542] and  $(\hat{u} - \phi)(\hat{x}) = 0$ . Then, there exist sequences  $\underline{\Omega}_k$  and  $\underline{x}_k \in \underline{\Omega}_k$  such that as  $k \rightarrow \infty$ ,

$$\begin{aligned} \hat{h}(\underline{\Omega}_k) &\rightarrow 0, \quad \underline{x}_k \rightarrow \hat{x}, \quad u^{\Omega_k}(\underline{x}_k) \rightarrow \hat{u}(\hat{x}), \quad \text{and} \\ (u^{\Omega_k} - \phi)(\underline{x}_k) &\geq (u^{\Omega_k} - \phi)(\underline{x}'_k) \text{ for all } \underline{x}'_k \in \mathcal{N}(\underline{x}_k). \end{aligned}$$

Let  $\xi_k = (u^{\Omega_k} - \phi)(\underline{x}_k)$ . We have  $\xi_k \rightarrow 0$  and  $\phi(\underline{x}'_k) + \xi_k \geq u^{\Omega_k}(\underline{x}'_k)$ , for all  $\underline{x}'_k \in \mathcal{N}(\underline{x}_k)$ . Consequently, by the monotonicity of  $\underline{H}$  (Theorem 3.6) and the definition of  $u^{\Omega_k}$  we have

$$\begin{aligned} \underline{H}(N(\underline{x}_k), \phi + \xi_k, \phi(\underline{x}_k) + \xi_k) &= \underline{H}(N(\underline{x}_k), \phi + \xi_k, u^{\Omega_k}(\underline{x}_k)) \\ &\leq \underline{H}(N(\underline{x}_k), u^{\Omega_k}, u^{\Omega_k}(\underline{x}_k)) = 0 \end{aligned}$$

Take the limit as  $k \rightarrow \infty$ , and use the consistency of the  $\underline{H}$  (Theorem 4.1) to get

$$\begin{aligned} 0 &\geq \lim_{k \rightarrow \infty} \underline{H}(N(\underline{x}_k), \phi + \xi_k, \phi(\underline{x}_k) + \xi_k) \\ &= \lim_{\substack{\underline{x}_0 \rightarrow x, \xi \rightarrow 0, \\ h_j^{s_j} \rightarrow 0, 1 \leq j \leq d}} \underline{H}(N(\underline{x}_0), \phi + \xi, \phi(\underline{x}_0) + \xi) = H(x, D\phi(x)) \end{aligned}$$

Therefore,  $\hat{u}$  is a viscosity subsolution.  $\square$

**REMARK 2.** We note that if  $H$  is continuous, (1.1) satisfies a comparison principal [4]: for any bounded upper semi-continuous  $u^*$  and bounded lower semi-continuous  $u_*$ , which are a viscosity subsolution and supersolution, respectively, of (1.1), such that  $u^* \leq u_*$  on  $\partial\Omega$ , we have  $u^* \leq u_*$  in  $\Omega$ .

**THEOREM 4.5.** Let  $H$  be continuous. The function  $\hat{u} = \check{u} = u$  is the unique viscosity solution of (1.1). As  $\hat{h}(\Omega) \rightarrow 0$ ,  $u^{\Omega}$  converges uniformly to  $u$ .

*Proof.* By Theorem 4.4,  $\hat{u}$  is an upper semi-continuous viscosity subsolution of (1.1) and  $\check{u}$  is a lower semi-continuous viscosity supersolution of (1.1). It follows by the comparison principal that  $\hat{u} \leq \check{u}$ . But  $\check{u} \leq \hat{u}$  by (4.5), so  $u = \hat{u} = \check{u}$  is a viscosity solution of (1.1). Again, by the comparison principal,  $u$  must be the unique viscosity solution of (1.1). Therefore, by (4.5)  $u^{\Omega}$  converges uniformly to  $u$ , as  $\hat{h}(\Omega) \rightarrow 0$ .  $\square$

**5. Experiments.** We conduct experiments to show numerical evidence that the result of FMM converges to the viscosity solution of (1.1), to demonstrate types of anisotropic problems that can be solved, and to determine the effectiveness of the node and simplex elimination techniques described in Appendix A. Throughout this section, the boundary conditions are  $g(x) = 0$  for  $x \in \partial\Omega$ . For all experiments below, excluding that in Section 5.4, we discretize  $[-1, 1]^d$  such that there are  $m$  uniformly-spaced nodes in each dimension and we ensure that there is a node at the origin  $O$ .

**5.1. Convergence Study.** We examine the difference between the solution to (3.3) and the solution to (1.1) for two simple Dirichlet problems. In particular, we look at how the absolute error changes as the grid spacing decreases toward zero. For the problems considered,  $\Omega = [-1, 1]^d \setminus \{O\}$ . We take  $H$  to have the form in (2.2), where  $G(Du(x)) = \|Du(x)\|_p$  and  $p = 1$  or  $p = 2$ . The boundary conditions are  $g(O) = 0$ . We use the analytic node value update equations provided in Appendix B.

Since there is a node at  $O$ , any error introduced is from the discretization of  $H$  and not from the discretization of the boundary condition. The approximation errors are summarized in Table 5.1.

**5.2. Asymmetric Anisotropic Problem.** For this anisotropic problem,  $H$  is as in (2.2), where  $G$  is defined by (2.5) (see Figure 2.3(b)). The domain is given by  $\Omega = [-1, 1]^2 \setminus \{O\}$  and  $\partial\Omega = \{O\}$ . The cost is  $c(x) = 1$ , except in four rectangular regions shown in black in Figure 5.1, where  $c(x) \gg 1$ . In the `Update` function, we analytically computed to solution to (3.4) using the equations for updating from a

$d$	$m$	$n$	$h$	$p = 1$				$p = 2$			
				$e_\infty$	$r_\infty$	$e_1$	$r_1$	$e_\infty$	$r_\infty$	$e_1$	$r_1$
2	11	1.2e2	2.0e-1	2.2e-1		6.3e-2		1.2e-1		6.2e-2	
	21	4.4e2	1.0e-1	1.7e-1	.41	3.7e-2	.77	7.8e-2	.56	4.3e-2	.53
	41	1.7e3	5.0e-2	1.2e-1	.46	2.0e-2	.85	5.0e-2	.65	2.8e-2	.63
	81	6.6e3	2.5e-2	8.8e-2	.48	1.1e-2	.90	3.1e-2	.70	1.7e-2	.69
	161	2.6e4	1.3e-2	6.3e-2	.49	5.7e-3	.94	1.8e-2	.75	1.0e-2	.73
	321	1.0e5	6.3e-3	4.4e-2	.49	2.9e-3	.96	1.1e-2	.78	6.1e-3	.77
	641	4.1e5	3.1e-3	3.1e-2	.50	1.5e-3	.97	6.1e-3	.81	3.5e-3	.79
	1281	1.6e6	1.6e-3	2.2e-2	.50	7.6e-4	.98	3.4e-3	.83	2.0e-3	.82
3	11	1.3e3	2.0e-1	3.5e-1		1.2e-1		2.1e-1		1.2e-1	
	21	9.3e3	1.0e-1	2.6e-1	.43	6.9e-2	.78	1.4e-1	.58	8.4e-2	.57
	41	6.9e4	5.0e-2	1.9e-1	.47	3.9e-2	.85	8.7e-2	.66	5.4e-2	.65
	81	5.3e5	2.5e-2	1.3e-1	.49	2.1e-2	.89	5.3e-2	.72	3.3e-2	.70
	161	4.2e6	1.3e-2	9.5e-2	.50	1.1e-2	.92	3.1e-2	.76	2.0e-2	.74
4	11	1.5e4	2.0e-1	4.4e-1		1.7e-1		2.9e-1		1.8e-1	
	21	1.9e5	1.0e-1	3.2e-1	.45	9.8e-2	.78	1.9e-1	.60	1.2e-1	.58
	41	2.8e6	5.0e-2	2.3e-1	.48	5.5e-2	.83	1.2e-1	.67	7.7e-2	.66

TABLE 5.1

Errors of approximate solution computed by FMM compared to exact solution of (1.1), where  $H$  is as in (2.2) and  $G(Du(x)) = \|Du(x)\|_p$ . The variables  $d$ ,  $m$ , and  $n$  are the dimension, the number of nodes in each dimension, and the total number of nodes, respectively. Other variables are the spacing  $h$  between grid nodes, the  $\mathcal{L}_\infty$ -error  $e_\infty$ , the  $\mathcal{L}_\infty$  convergence rate  $r_\infty$ , the  $\mathcal{L}_1$ -error  $e_1$ , and the  $\mathcal{L}_1$  convergence rate  $r_1$ .




FIG. 5.1. Contours of  $\underline{u}$  computed for the anisotropic problem where Hamiltonian  $H$  is as in (2.2) and  $G$  is as in (2.5). The black circle at  $O = (0, 0)$  indicates  $\partial\Omega$  and in the black rectangles,  $c(x) \gg 1$ . In these regions,  $\underline{u}$  has purposefully not been computed.

single simplex given in Appendix B. The number of nodes in each dimension is  $m = 1281$ . We plot the contours of  $\underline{u}$  computed by FMM in Figure 5.1. Note the asymmetric contours where the characteristics bend through gaps. The relationship between the shape of the contours of  $G$  in Figure 2.3(b) and those of  $\underline{u}$  is explained by the duality articulated in Proposition 2.7 of [2].




FIG. 5.2. Using FMM for computation of travel times of qP-waves in two-dimensional VTI media. (a) Contours of Hamiltonian  $H$  as given in (5.1). (b) Contours of approximate solution  $\underline{u}$  computed by FMM.

**5.3. Anelliptic Elastic Wave Propagation.** We consider elastic wave propagation in transversely isotropic media with a vertical axis of symmetry (VTI media) as is done in [12]. In particular, we wish to find the arrival times of qP-waves propagating in two dimensions from a point source at the origin  $O$ . We solve the anisotropic HJ PDE given by defining the Hamiltonian

$$H(q) = \frac{1}{2}(q_1^2 + q_2^2) \left\{ (a+l)q_1^2 + (c+l)q_2^2 + \sqrt{[(a-l)q_1^2 - (c-l)q_2^2]^2 + 4(f+l)^2 q_1^2 q_2^2} \right\} - 1. \quad (5.1)$$

This Hamiltonian is derived from the anisotropic Eikonal equation and the exact qP-wave phase velocity equation in [12]. The parameters  $a = 14.47$ ,  $l = 2.28$ ,  $c = 9.57$ , and  $f = 4.51$  are taken from [12].

The Hamiltonian  $H$  and the approximate solution  $\underline{u}$  resulting from FMM are shown in Figure 5.2. We have not shown analytically that (5.1) satisfies strict one-sided monotonicity for some range of parameters. However, the level sets of  $H$  as shown in Figure 5.2(a) indicate that  $H$  is strictly one-sided monotone for the given parameters. Furthermore, the level sets of  $H$  indicate the  $H$  is convex and computation of the derivative of  $H$  using the symbolic mathematics program Maple shows that  $H$  satisfies Osher's criterion for the given parameters. As a result, the analysis in this paper can be applied to the problem and FMM can be used to compute the solution.

We used a grid of size  $201 \times 201$ . In the `Update` function, we used the interval method to solve (3.4) numerically. We computed the maximum relative error of  $\underline{u}$  to be 0.0076, when compared to the travel-time computed with the group-velocity approximation for qP-waves presented in [12]. In turn, the group-velocity approximation is claimed to have a maximum error of about 0.003, when compared to the true solution.

**5.4. Two Robots.** We consider the two-robot coordinated navigation problem illustrated in Figure 5.3. The circular robots are free to move independently in a 2-dimensional plane but may not collide with each other or the obstacles (black region). Each may travel at a maximum speed of  $1/c(x)$  in any direction. The robots attempt to achieve a joint goal state. This goal should be achieved in minimal time from any initial state in the domain without incurring collisions.

Let the state of the dark-colored robot be  $(x_1, x_2) \in \mathbb{R}^2$  and the state of the light-colored robot be  $(x_3, x_4) \in \mathbb{R}^2$  so that the combined state of the two robots is  $(x_1, x_2, x_3, x_4) \in \mathbb{R}^4$ . We define the control-theoretic action set

$$\mathcal{A}(x) = \{a \mid F(a) = \|(\|(a_1, a_2)\|_2, \|(a_3, a_4)\|_2)\|_\infty \leq 1/c(x)\}.$$

Proposition 2.7 of [2] states that we can use the dual of  $F$  to obtain

$$G(x, Du(x)) = \|(\|(\partial_1 u(x), \partial_2 u(x))\|_2, \|(\partial_3 u(x), \partial_4 u(x))\|_2)\|_1, \quad (5.2)$$

where  $Du(x) = (\partial_1 u(x), \partial_2 u(x), \partial_3 u(x), \partial_4 u(x))$ . Where  $x$  is a collision state, we set  $c(x) \gg 1$ . For all other states  $x$ ,  $c(x) = 1$ .

We can compute  $\underline{u}$  using FMM since  $G$  is a mixed  $p$ -norm, and thus  $H$  satisfies Properties 1 to 4 (see Section 2.2). The domain  $\Omega$  is discretized using a uniform orthogonal grid of  $(81 \times 21)^2$  nodes. The discretization of (5.2) is quartic in  $\underline{u}_0$ , so it is difficult to solve analytically. However, Theorem 3.5 tells us that we can determine the solution to (3.4) uniquely. As a result, numerical root-finders can easily be used to compute this solution in the `Update` function.

Once we have determined  $\underline{u}$  using FMM, we approximately solve the following ODE to determine an optimal collision-free trajectory from any initial state to the goal:

$$\frac{dx}{dt} = \operatorname{argmax}_{a \in \mathcal{A}(x)} (-D\underline{u}(x) \cdot a) - 1. \quad (5.3)$$

Solving this ODE is not the focus of the paper so we take a simple approach. The ODE (5.3) is discretized using forward Euler. The gradient  $D\underline{u}(x)$  is determined by a first-order finite difference scheme. At each time step, each robot moves at its maximum speed  $1/c(x)$  in the direction of the relevant components of the negative gradient, e.g. (for dark-colored robot):

$$(a_1, a_2) = \frac{-1}{c(x)} \frac{(\partial_1 \underline{u}(x), \partial_2 \underline{u}(x))}{\|(\partial_1 \underline{u}(x), \partial_2 \underline{u}(x))\|}.$$

If the relevant components of the gradient fall below a small threshold the robot will not move at all, as is the case for the light-colored robot in Figures 5.3(d) and 5.3(e). The optimal trajectories from a single starting condition are shown in Figure 5.3.

**5.5. Efficient Implementation.** Appendix A describes three different methods for improving the efficiency of the `Update` function: symmetry, causality, and solution elimination. Some of these methods are related to those found in [17, 33]. However, experimental results indicate that the efficiency gains from using these methods are not substantial for an already efficient implementation of FMM. In such an implementation the `Update` function only computes updates from those nodes that have already been extracted from  $\mathcal{Q}$  using the `ExtractMin` function. Also, only simplices that include the most-recently extracted node  $\underline{y}$  are considered in `Update`. Our experiments show that in many calls to `Update`, only a single simplex fits these criteria,






FIG. 5.3. *Two-robot coordinated optimal navigation problem. The joint goal is for the dark-colored robot to reach the center of the upper bulb and light-colored robot to reach the center of the lower bulb. Black indicates an obstacle region. The sequence shows the robots achieving their joint goal without collision from a particular initial state. The solution of (1.1), where  $H$  is given by (2.2) and  $G$  is given by (5.2) allows quick determination of the optimal collision-free trajectories for both robots from any initial condition [1].*

and the fraction of updates for which only a single simplex fits the criteria grows as the grid is refined. For this reason, further techniques for eliminating nodes and simplices, such as those described in Appendix A, are largely ineffective.

However, for coarse grid resolutions and problems where characteristics intersect often, multiple simplices are considered by `Update` frequently enough that symmetry elimination, which is very cheap, significantly improves efficiency. In some cases, a node value update can be ignored altogether if the most-recently extracted node is eliminated by symmetry.

Despite the fact that the node and simplex elimination techniques described in Appendix A are useful only in limited circumstances we include them for theoretical interest and because they may be applied in other algorithms, such as sweeping methods, that also require the `Update` function.

**6. Conclusion.** We have described a new class of static HJ PDEs with axis-aligned but potentially asymmetric anisotropy. Assuming Properties 1 to 4 of the Hamiltonian, we showed that uniqueness, monotonicity and causality hold for a standard finite difference discretization of these PDEs on an orthogonal grid, and so the Fast Marching Method can be used to approximate their solution. In the appendix, we also demonstrate several methods for reducing the number of neighboring simplices which must be considered when computing node updates, including novel methods which work when the PDE and/or grid are asymmetric. In future work, these results might be generalized to unstructured grids.

**Acknowledgments.** We would like to thank Alexander Vladimirsky for enlightening discussions about FMM and OUM, and Adam Oberman for discussions and insights regarding [5].

### Appendix A. Efficient Implementation of Update.

We discuss ways to improve the efficiency of the `Update` function, which calculates the unique solution  $\mu = \tilde{\mu}$  to (3.4). We note that these improvements may be used for any type of solution method, including FMM and sweeping methods, as long as (3.4) is being solved. Some efficiency improvements are related to similar ideas specific to the isotropic Eikonal equation found in [17, 33].

Efficiency can be gained by determining which neighbors  $\underline{x} \in \mathcal{N}(\underline{x}_0)$  have no influence on the solution and eliminating them from consideration. Let

$$\sigma = (\sigma_1, \sigma_2, \dots, \sigma_d),$$

where  $\sigma_j \subseteq \{\pm 1\}$ , indicate which  $\underline{x} \in \mathcal{N}$  are considered in determining the solution  $\mu = \tilde{\mu}$ . Let  $\mathcal{N}_\sigma$  be the reduced set of neighbor nodes defined by  $\sigma$ . Let  $\mathcal{S}_\sigma$  be the set of neighboring simplices that can be formed by the neighbors in  $\mathcal{N}_\sigma$ . For example, in  $d = 4$  dimensions, take

$$\sigma = (\emptyset, \{\pm 1\}, \{-1\}, \{\pm 1\}).$$

We have

$$\begin{aligned} \mathcal{N}_\sigma &= \{\underline{x}_2^\pm, \underline{x}_3^-, \underline{x}_4^\pm\} \quad \text{and} \\ \mathcal{S}_\sigma &= \{(0, -1, -1, -1), (0, +1, -1, -1), (0, -1, -1, +1), (0, +1, -1, +1)\}. \end{aligned}$$

Let  $\underline{H}_\sigma(N, \phi, \mu) = \max_{s \in \mathcal{S}_\sigma} [H(\underline{x}_0, \underline{D}^s(N, \phi, \mu))]$  be the *reduced-neighbor numerical Hamiltonian*, a modification of (3.1) that only considers the neighbors and simplices indicated by  $\sigma$ . For  $s \in \mathcal{S}_\sigma$  and  $1 \leq j \leq d$ ,  $s_j = 0$  indicates that  $\underline{x}_j^{s_j}$  is not considered in computing the gradient approximation  $D^s \underline{u}(\mu)$ ; that is,  $\underline{D}_j^s(N, \phi, \mu) = 0$  if  $s_j = 0$  and  $\underline{D}_j^s$  satisfies (3.2) otherwise.

To implement `Update` we first reduce the set of considered neighbors and then solve

$$\underline{H}_\sigma(N(\underline{x}_0), \underline{u}, \mu) = 0 \tag{A.1}$$

for  $\mu = \tilde{\mu}$  to determine a node's value  $\underline{u}(\underline{x}_0)$ . As in Section 3, we may write  $\underline{H}_\sigma(\mu) = \underline{H}_\sigma(N, \phi, \mu)$  and  $\underline{D}^s(\mu) = \underline{D}^s(N, \phi, \mu)$ , where no ambiguity results. Note that some properties of (A.1) are retained from (3.4) as long as at least one considered neighbor remains in  $\sigma$ . Let

$$\check{\underline{u}}_\sigma = \min_{\underline{x} \in \mathcal{N}_\sigma} (\underline{u}(\underline{x})).$$

PROPOSITION A.1. (analogue of Lemma 3.3)  $\underline{H}_\sigma(\mu)$  is strictly increasing on  $\mu \geq \check{\underline{u}}_\sigma$ .

PROPOSITION A.2. (analogue of Lemma 3.4) The numerical Hamiltonian  $\underline{H}_\sigma(\mu)$  satisfies the following.

- (a)  $\underline{H}_\sigma(\mu) = H(0) < 0$  for  $\mu \leq \check{\underline{u}}_\sigma$ .
- (b)  $\underline{H}_\sigma(\mu) \rightarrow \infty$  as  $\mu \rightarrow \infty$ .
- (c)  $\underline{H}_\sigma(\mu)$  is nondecreasing on all  $\mu$ .

PROPOSITION A.3. (analogue of Theorem 3.5) There exists a unique solution  $\mu = \tilde{\mu}$  to  $\underline{H}_\sigma(\mu) = 0$  such that  $\tilde{\mu} > \check{\underline{u}}_\sigma$ .

**A.1. Symmetry.** We show how the considered neighbors  $\sigma$  can be reduced by keeping only the neighbor with smaller value of a pair of opposite neighbors in the  $j$ th dimension when (3.1) is symmetric in that dimension. This procedure is a generalization of those in [17, 33] to all axis-aligned anisotropic problems on unequally spaced grids. First, we introduce useful notation.

Let  $q \in \mathbb{R}^d$ . Let  $T^i(q)$  be a reflection of  $q$  in the hyperplane orthogonal to the  $i$ th axis, such that

$$T_j^i(q) = \begin{cases} -q_j, & \text{if } j = i, \\ q_j, & \text{otherwise,} \end{cases}$$

for  $1 \leq j \leq d$ . Let  $\Psi_j$  indicate symmetry of (3.1) in the  $j$ th dimension, as follows:

$$\Psi_j = \begin{cases} 1, & \text{if } |h_j^-| = |h_j^+| \text{ and for all } q \in \mathbb{R}^d, H(q) = H(T^j(q)), \\ 0, & \text{otherwise.} \end{cases}$$

In other words,  $\Psi_j = 1$  if and only if the grid spacing and  $H$  are symmetric in the  $j$ th dimension.

LEMMA A.4. *Let  $j$  be such that  $1 \leq j \leq d$ . Let  $s \in \mathcal{S}$ . Let  $s' = T^j(s)$ . If  $\Psi_j = 1$  and  $\underline{u}_j^{s'} \leq \underline{u}_j^s$ , then  $G(D^{s'}\underline{u}(\mu)) \geq G(D^s\underline{u}(\mu))$ , for all  $\mu$ .*

*Proof.* Let  $j$ ,  $s$ , and  $s'$  be as defined above. Let  $\Psi_j = 1$  and  $\underline{u}_j^{s'} \leq \underline{u}_j^s$ . Consider the components of  $T^j(D^{s'}\underline{u}(\mu))$ . We have

$$\begin{aligned} & T_j^j(D^{s'}\underline{u}(\mu))D_j^s\underline{u}(\mu) \\ &= -D_j^{s'}\underline{u}(\mu)D_j^s\underline{u}(\mu) \\ &= -\frac{\max(0, \mu - \underline{u}_j^{s'})}{-h_j^{s'}} \frac{\max(0, \mu - \underline{u}_j^s)}{-h_j^s} \geq 0, \end{aligned}$$

since  $h_j^{s'} h_j^s < 0$ . Furthermore,

$$\begin{aligned} |T_j^j(D^{s'}\underline{u}(\mu))| &= | -D_j^{s'}\underline{u}(\mu) | \\ &= \left| \frac{\max(0, \mu - \underline{u}_j^{s'})}{-h_j^{s'}} \right| \\ &\geq \left| \frac{\max(0, \mu - \underline{u}_j^s)}{-h_j^s} \right| = |D_j^s\underline{u}(\mu)|, \end{aligned}$$

since  $h_j^{s'} = -h_j^s$  and  $\underline{u}_j^{s'} \leq \underline{u}_j^s$ . For  $i \neq j$ ,

$$T_i^j(D^{s'}\underline{u}(\mu)) = D_i^{s'}\underline{u}(\mu) = D_i^s\underline{u}(\mu),$$

since  $s'_i = s_i$ . Consequently,  $T^j(D^{s'}\underline{u}(\mu)) \supseteq D^s\underline{u}(\mu)$ .

Therefore, by the symmetry of  $G$  in the  $j$ th dimension and by the one-sided monotonicity of  $G$ ,

$$G(D^{s'}\underline{u}(\mu)) = G(T^j(D^{s'}\underline{u}(\mu))) \geq G(D^s\underline{u}(\mu)).$$

□

THEOREM A.5. Let  $\sigma$  be such that  $\sigma_j \subseteq \{\pm 1\}$ , for  $1 \leq j \leq d$ . Let  $\tilde{\sigma}$  be defined by

$$\tilde{\sigma}_j = \begin{cases} \{-1\}, & \text{if } \sigma_j = \{\pm 1\}, \Psi_j = 1 \text{ and } \underline{u}_j^- \leq \underline{u}_j^+, \\ \{+1\}, & \text{if } \sigma_j = \{\pm 1\}, \Psi_j = 1 \text{ and } \underline{u}_j^- > \underline{u}_j^+, \\ \sigma_j, & \text{otherwise,} \end{cases}$$

for  $1 \leq j \leq d$ . Let  $\mu = \mu_\sigma$  be the unique solution to  $\underline{H}_\sigma(\mu) = 0$ . Let  $\mu = \mu_{\tilde{\sigma}}$  be the unique solution to  $\underline{H}_{\tilde{\sigma}}(\mu) = 0$ . Then  $\mu_{\tilde{\sigma}} = \mu_\sigma$ .

*Proof.* Let  $\sigma, \tilde{\sigma}, \mu_\sigma$  and  $\mu_{\tilde{\sigma}}$  be as defined above. Consider a sequence  $\sigma^0, \sigma^1, \dots, \sigma^d$ , such that

$$\sigma_j^i = \begin{cases} \tilde{\sigma}_j^i, & \text{if } i \geq j, \\ \sigma_j^i, & \text{otherwise,} \end{cases}$$

for  $1 \leq j \leq d$ . Let  $\mu = \mu_{\sigma^i}$  be the unique solution to  $\underline{H}_{\sigma^i}(\mu) = 0$ .

Let  $i$  be such that  $1 \leq i \leq d$ . Assume  $\mu_{\sigma^{i-1}} = \mu_\sigma$ . For each  $s \in \mathcal{S}_{\sigma^{i-1}}$ , if  $s \notin \mathcal{S}_{\sigma^i}$ , then  $s' = T^i(s) \in \mathcal{S}_{\sigma^i}$ ,  $\Psi_i = 1$ , and  $\underline{u}_i^{s'} \leq \underline{u}_i^{s_i}$ . So, by Lemma A.4,  $G(D^{s'}\underline{u}(\mu)) \geq G(D^s\underline{u}(\mu))$ , for all  $\mu$ . In particular,  $G(D^{s'}\underline{u}(\mu_{\sigma^{i-1}})) \geq G(D^s\underline{u}(\mu_{\sigma^{i-1}}))$ . As a result, for each  $s \in \mathcal{S}_{\sigma^{i-1}}$ , there exists  $\tilde{s} \in \mathcal{S}_{\sigma^i}$ , such that  $G(D^{\tilde{s}}\underline{u}(\mu_{\sigma^{i-1}})) \geq G(D^s\underline{u}(\mu_{\sigma^{i-1}}))$ . Also, note that  $\mathcal{S}_{\sigma^i} \subseteq \mathcal{S}_{\sigma^{i-1}}$ . Consequently,

$$\begin{aligned} \underline{H}_{\sigma^i}(\mu_{\sigma^{i-1}}) &= \max_{s \in \mathcal{S}_{\sigma^i}} [G(D^s\underline{u}(\mu_{\sigma^{i-1}}))] - c \\ &= \max_{s \in \mathcal{S}_{\sigma^{i-1}}} [G(D^s\underline{u}(\mu_{\sigma^{i-1}}))] - c = \underline{H}_{\sigma^i}(\mu_{\sigma^{i-1}}) = 0, \end{aligned}$$

Accordingly  $\mu = \mu_{\sigma^{i-1}}$  is the unique solution to  $\underline{H}_{\sigma^i}(\mu) = 0$ . By the definition of  $\mu_{\sigma^i}$  and the assumption,  $\mu_{\sigma^i} = \mu_{\sigma^{i-1}} = \mu_\sigma$ . Therefore, by induction on  $i$ ,  $\mu_{\tilde{\sigma}} = \mu_\sigma$ , since  $\sigma^0 = \sigma$  and  $\sigma^d = \tilde{\sigma}$ . □

An implementation of the **Update** function can use the result obtained in Theorem A.5 to eliminate  $\underline{x} \in \mathcal{N}$  from consideration in solving (A.1) by exploiting symmetries in (3.1). We call this *symmetry elimination*.

REMARK 3. Theorem A.5 can be generalized to an asymmetric version. We let  $1 \leq j \leq d$  and let  $s_j, \tilde{s}_j \in \{\pm 1\}$  such that  $s_j \neq \tilde{s}_j$ . Node  $\underline{x}_j^{s_j} \in \mathcal{N}$  may be eliminated from consideration if

- $|h_j^{\tilde{s}_j}| \leq |h_j^{s_j}|$ ;
- for all  $q \in \mathbb{R}^d$  such that  $s_j q_j \geq 0$ ,  $H(q) \leq H(T^j(q))$ ;
- and  $\underline{u}_j^{\tilde{s}_j} \leq \underline{u}_j^{s_j}$ .

**A.2. Causality.** The causality of (3.1) can also be exploited to eliminate  $\underline{x} \in \mathcal{N}_\sigma$  from consideration. This observation was used in two distinct but equivalent methods for analytically computing the **Update** from a single simplex to solve an isotropic Eikonal equation [17, 33]. We show with the following theorem that the condition  $\underline{H}_\sigma(\underline{u}(\underline{x})) \geq 0$  can be checked to determine that a node  $\underline{x}$  is non-causal, i.e., that the solution  $\mu = \mu_\sigma$  to (A.1) is not dependent on the node  $\underline{x}$  and its value  $\underline{u}(\underline{x})$ .

THEOREM A.6. Let  $\sigma$  be such that  $\sigma_j \subseteq \{\pm 1\}$ , for  $1 \leq j \leq d$ . Pick any  $s \in \mathcal{S}_\sigma$  and  $i \in \{1, 2, \dots, d\}$ , such that  $s_i \neq 0$  and  $\underline{H}_\sigma(\underline{u}_i^{s_i}) \geq 0$ . Let  $\tilde{\sigma}$  be defined by

$$\tilde{\sigma}_j = \begin{cases} \sigma_j \setminus \{s_j\}, & \text{if } j = i, \\ \sigma_j, & \text{otherwise,} \end{cases}$$

Let  $\mu = \mu_\sigma$  be the unique solution to  $\underline{H}_\sigma(\mu) = 0$ . Let  $\mu = \mu_{\tilde{\sigma}}$  be the unique solution to  $\underline{H}_{\tilde{\sigma}}(\mu) = 0$ . Then  $\mu_{\tilde{\sigma}} = \mu_\sigma$ .

*Proof.* Let  $\sigma, s, i, \tilde{\sigma}, \mu_\sigma$  and  $\mu_{\tilde{\sigma}}$  be as defined above. By Proposition A.2(c),  $\underline{H}_\sigma(\mu)$  is nondecreasing. Since  $\underline{H}_\sigma(\underline{u}_i^{s_i}) \geq 0 = \underline{H}_\sigma(\mu_\sigma)$ , it must be that  $\mu_\sigma \leq \underline{u}_i^{s_i}$ . Note that  $\underline{H}_{\tilde{\sigma}}(\mu)$  is identical to  $\underline{H}_\sigma(\mu)$  except for  $D_i^s \underline{u}(\mu)$  which is set to zero in  $\underline{H}_{\tilde{\sigma}}(\mu)$ . But for  $\mu \leq \underline{u}_i^{s_i}$ , we also have  $D_i^s \underline{u}(\mu) = 0$  in  $\underline{H}_\sigma(\mu)$ . Consequently,  $\underline{H}_{\tilde{\sigma}}(\mu) = \underline{H}_\sigma(\mu)$  for  $\mu \leq \underline{u}_i^{s_i}$ . In particular,  $\underline{H}_{\tilde{\sigma}}(\mu_\sigma) = \underline{H}_\sigma(\mu_\sigma) = 0$ . Therefore,  $\mu_{\tilde{\sigma}} = \mu_\sigma$ .  $\square$

Theorem A.6 states that the unique solution  $\mu$  to (A.1) does not change when a non-causal node is removed from  $\sigma$ . This node removal can be repeated until all non-causal nodes have been removed and the solution  $\mu = \mu_\sigma$  will remain unchanged. We call this *causality elimination*. A binary or linear search through sorted neighbors' values can be used to determine the largest node value that might be causal. Note that causality elimination does not require symmetry in (3.1). However, the test for non-causality requires an evaluation of  $\underline{H}_\sigma$ , which is more expensive than the comparison of two neighbors' values used for symmetry elimination.

**A.3. Solution.** After eliminating from consideration nodes in  $\sigma$  using symmetry and causality elimination, we can determine the solution  $\mu = \tilde{\mu}$  to (A.1). Let

$$\tilde{\mu} = \min_{s \in \mathcal{S}_\sigma} (\mu_s), \quad (\text{A.2})$$

where  $\mu = \mu_s$  is the unique solution to

$$H(D^s \underline{u}(\mu)) = 0. \quad (\text{A.3})$$

We show with the following proposition that, instead of solving (A.1) directly, we can solve (A.3) for each  $s \in \mathcal{S}_\sigma$  and take the minimum such solution  $\tilde{\mu}$ . It can be shown that  $H(D^s \underline{u}(\mu))$  is continuous and nondecreasing on  $\mu$  and that (A.3) has a unique solution in an analogous but simpler manner as the proof of Theorem 3.5.

**PROPOSITION A.7.** *Let  $\hat{\mu}$  be the unique solution to (A.1). Then  $\hat{\mu} = \tilde{\mu}$ .*

*Proof.* Let  $\mu_s, \tilde{\mu}$  and  $\hat{\mu}$  be as defined above. For any  $s \in \mathcal{S}_\sigma$ , we know  $\mu_s \geq \tilde{\mu}$ . Since  $H(D^s \underline{u}(\mu))$  is nondecreasing on  $\mu$ , it must be that  $H(D^s \underline{u}(\mu)) \leq H(D^s \underline{u}(\mu_s)) = 0$ , for all  $\mu \leq \mu_s$ . In particular,  $H(D^s \underline{u}(\tilde{\mu})) \leq 0$ . Furthermore, by the definition of  $\tilde{\mu}$ , there exists an  $\tilde{s} \in \mathcal{S}_\sigma$ , such that  $H(D^{\tilde{s}} \underline{u}(\tilde{\mu})) = 0$ . Consequently,

$$\underline{H}_\sigma(\tilde{\mu}) = \max_{s \in \mathcal{S}_\sigma} H(D^s \underline{u}(\tilde{\mu})) = 0. \quad (\text{A.4})$$

Therefore,  $\hat{\mu} = \tilde{\mu}$  solves (A.1) and it is a unique solution by Proposition A.3.  $\square$

We further show that we may be able to determine  $\tilde{\mu}$  without solving (A.3) for each  $s \in \mathcal{S}_\sigma$ . We demonstrate using the following proposition that if we have computed a solution  $\mu = \mu_s$  of (A.3) for some  $s \in \mathcal{S}_\sigma$ , we can easily determine if  $\mu_{\tilde{s}} \geq \mu_s$ , where  $\mu = \mu_{\tilde{s}}$  is the solution to  $H(D^{\tilde{s}} \underline{u}(\mu)) = 0$  for some other  $\tilde{s} \in \mathcal{S}_\sigma$ . Note we do not necessarily need to compute  $\mu_{\tilde{s}}$  to rule it out as a minimal solution.

**PROPOSITION A.8.** *Let  $s \in \mathcal{S}_\sigma$  and  $\tilde{s} \in \mathcal{S}_\sigma$ . Let  $\mu = \mu_s$  be the unique solution to  $H(D^s \underline{u}(\mu)) = 0$  and  $\mu = \mu_{\tilde{s}}$  be the unique solution to  $H(D^{\tilde{s}} \underline{u}(\mu)) = 0$ . Then  $\mu_{\tilde{s}} < \mu_s$  if and only if  $H(D^{\tilde{s}} \underline{u}(\mu_s)) > H(D^s \underline{u}(\mu_s))$ .*

*Proof.* Let  $\mu_s$  and  $\mu_{\tilde{s}}$  be as defined above. If  $H(D^{\tilde{s}} \underline{u}(\mu_s)) > H(D^s \underline{u}(\mu_s)) = 0$ , then the unique solution  $\mu = \mu_{\tilde{s}}$  to  $H(D^{\tilde{s}} \underline{u}(\mu)) = 0$  must be such that  $\mu_{\tilde{s}} < \mu_s$ , since  $H(D^{\tilde{s}} \underline{u}(\mu))$  is nondecreasing on  $\mu$ . Similarly, if  $H(D^{\tilde{s}} \underline{u}(\mu_s)) \leq H(D^s \underline{u}(\mu_s))$  then the unique solution  $\mu = \mu_{\tilde{s}}$  to  $H(D^{\tilde{s}} \underline{u}(\mu)) = 0$  must be such that  $\mu_{\tilde{s}} \geq \mu_s$ .  $\square$

The result of Proposition A.8 can be used to eliminate simplices  $s \in \mathcal{S}_\sigma$  for which solutions to (A.3) are irrelevant to the computation. We call this process *solution elimination*.

### Appendix B. Analytic Solutions.

We provide analytic node value update equations for the cases where  $H$  is given by (2.2), where  $G(Du(x)) = \|Du(x)\|_p$  and  $p = 1$ ,  $p = 2$ , or  $p = \infty$ . In these cases, there is an exact solution to (3.4). The equation for  $p = 2$  fixes some errors in the appendix of [18]. In [1] we demonstrated that these cases could be treated by FMM and are useful for robotic applications. However, here we generalize the update equations to any dimension and grid spacing.

Let  $(v_1, v_2, \dots, v_m)$  be the values of the neighboring nodes in the simplex  $s \in \mathcal{S}_\sigma$  and  $(h_1, h_2, \dots, h_m)$  be the corresponding grid spacings. We are solving for  $\mu$ . In order to use the analytic updates below, non-causal node values must already have been eliminated using causality elimination, so  $\mu > \max_{1 \leq j \leq m} v_j$ . However, in the case of the efficient implementation of FMM discussed in Section 5.5, any nodes that would be removed from consideration by causality elimination could not already have been extracted from  $\mathcal{Q}$ , and so the analytic updates below can be applied directly.

**B.1. Update for  $p = 1$ .** From (A.3) we have

$$\sum_j \left( \frac{|\mu - v_j|}{h_j} \right) = c.$$

Assume  $\mu > \max_{1 \leq j \leq m} v_j$  and multiply through by  $\prod_l h_l$  to obtain

$$\sum_j \left( \prod_{l \neq j} h_l \right) \mu - \sum_j \left( \prod_{l \neq j} h_l \right) v_j = \left( \prod_l h_l \right) c.$$

Then solve for  $\mu$  to get

$$\mu = \frac{\sum_j \left( \prod_{l \neq j} h_l \right) v_j + \prod_l h_l c}{\sum_j \prod_{l \neq j} h_l}.$$

**B.2. Update for  $p = 2$ .** From (A.3) we have

$$\sum_j \left( \frac{\mu - v_j}{h_j} \right)^2 = c^2.$$

Multiply through by  $\prod_l h_l^2$  to get

$$\left[ \sum_j \prod_{l \neq j} h_l^2 \right] \mu^2 - 2 \left[ \sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j \right] \mu + \sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j^2 - \left( \prod_l h_l^2 \right) c^2 = 0.$$

Then, using the quadratic formula, solve for  $\mu$ :

$$\begin{aligned} \mu &= \frac{2 \sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j + \sqrt{\left[ 2 \sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j \right]^2 - 4 \left[ \sum_j \prod_{l \neq j} h_l^2 \right] \left[ \sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j^2 - \left( \prod_l h_l^2 \right) c^2 \right]}}{2 \sum_j \prod_{l \neq j} h_l^2} \\ &= \frac{\sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j + \prod_l h_l \sqrt{\left( \sum_j \prod_{l \neq j} h_l^2 \right) c^2 + \sum_{j_1} \sum_{j_2} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1} v_{j_2} - \sum_{j_1} \sum_{j_2} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1}^2}}{\sum_j \prod_{l \neq j} h_l^2}. \end{aligned}$$

We only consider the larger of the two quadratic solutions since the alternative will result in  $\mu \leq \max_{1 \leq j \leq m} v_j$ . The last two terms of the discriminant can be made more concise as follows:

$$\begin{aligned} & \sum_{j_1} \sum_{j_2} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1} v_{j_2} - \sum_{j_1} \sum_{j_2} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1}^2 \\ &= \sum_{j_1} \sum_{j_2 \neq j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1} v_{j_2} + \sum_{j_1} \sum_{j_2 = j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1} v_{j_2} \\ & \quad - \sum_{j_1} \sum_{j_2 \neq j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1}^2 - \sum_{j_1} \sum_{j_2 = j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1}^2 \\ &= 2 \sum_{j_1} \sum_{j_2 > j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) v_{j_1} v_{j_2} + \sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j^2 \\ & \quad - \sum_{j_1} \sum_{j_2 > j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) (v_{j_1}^2 + v_{j_2}^2) - \sum_j \left( \prod_{l \neq j} h_l^2 \right) v_j^2 \\ &= \sum_{j_1} \sum_{j_2 > j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) 2v_{j_1} v_{j_2} - \sum_{j_1} \sum_{j_2 > j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) (v_{j_1}^2 + v_{j_2}^2) \\ &= - \sum_{j_1} \sum_{j_2 > j_1} \left( \prod_{l \neq j_1, j_2} h_l^2 \right) (v_{j_1} - v_{j_2})^2. \end{aligned}$$

**B.3. Update for  $p = \infty$ .** From (A.3) we have

$$\max_j \left( \frac{|\mu - v_j|}{h_j} \right) = c.$$

Assume  $\mu > \max_{1 \leq j \leq m} v_j$  solve for  $\mu$  to obtain

$$\mu = \min_j (v_j + h_j c).$$

The  $p = \infty$  case is identical to the update formula for Dijkstra’s algorithm for shortest path on a discrete graph.

#### REFERENCES

- [1] K. Alton and I. Mitchell. Optimal path planning under different norms in continuous state spaces. In *Proceedings of the International Conference on Robotics and Automation*, pages 866–872, 2006.
- [2] K. Alton and I. M. Mitchell. Fast marching methods for a class of anisotropic stationary Hamilton-Jacobi equations. Technical Report TR-2006-27, Department of Computer Science, University of British Columbia, 2007.
- [3] K. Alton and I. M. Mitchell. Fast marching methods for Hamilton-Jacobi equations with axis-aligned anisotropy. 2008. Accepted for publication in *SIAM J. NUMER. ANAL.*
- [4] M. Bardi and I. Capuzzo-Dolcetta. *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations*. Birkhauser, Boston and Basel and Berlin, 1997.
- [5] G. Barles and P. E. Souganidis. Convergence of approximation schemes for fully nonlinear second order equations. *Asymptot. Anal.*, 4:271–283, 1991.
- [6] F. Bornemann and C. Rasch. Finite-element discretization of static Hamilton-Jacobi equations based on a local variational principle. *Comput. Vis. Sci.*, 9:57–69, 2006.
- [7] M. Boue and P. Dupuis. Markov chain approximations for deterministic control problems with affine dynamics and quadratic cost in the control. *SIAM J. NUMER. ANAL.*, 36(3):667–695, 1999.
- [8] M. G. Crandall, H. Ishii, and P. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc.*, 27(1):1–67, 1992.
- [9] P.-E. Danielsson. Euclidean distance mapping. *Comput. Graphics Image Process.*, 14(3):227–248, Nov. 1980.
- [10] E. W. Dijkstra. A note on two problems in connection with graphs. *Numer. Math.*, 1(1):269–271, Dec. 1959.
- [11] L. C. Evans. *Partial Differential Equations*. American Mathematical Society, Providence, Rhode Island, 1991.
- [12] S. Fomel. On anelliptic approximations for qp velocities in vti media. *Geophysical Prospecting*, 52:247–259, April 2004.
- [13] P. A. Gremaud and C. M. Kuster. Computational study of fast methods for the Eikonal equation. *SIAM J. SCI. COMPUT.*, 27:1803–1816, 2006.
- [14] S.-R. Hysing and S. Turek. The Eikonal equation: Numerical efficiency vs. algorithmic complexity on quadrilateral grids. In *Proceedings of Algorithm 2005*, pages 22–31, 2005.
- [15] C. Y. Kao, S. Osher, and Y. Tsai. Fast sweeping methods for static Hamilton-Jacobi equations. *SIAM J. NUMER. ANAL.*, 42(6):2612–2632, 2004-2005.
- [16] R. Kimmel and J. A. Sethian. Fast marching methods on triangulated domains. *Proc. Natl. Acad. Sci.*, 95:8341–8435, 1998.
- [17] R. Kimmel and J. A. Sethian. Optimal algorithm for shape from shading and path planning. *J. Math. Imaging Vision*, 14(3):237–244, May 2001.
- [18] I. M. Mitchell and S. Sastry. Continuous path planning with multiple constraints. Technical Report UCB/ERL M03/34, UC Berkeley Engineering Research Laboratory, August 2003.
- [19] S. Osher and R. P. Fedkiw. Level set methods: An overview and some recent results. *J. Comput. Phys.*, 169(2):463–502, May 2001.
- [20] L. C. Polymenakos, D. P. Bertsekas, and J. N. Tsitsiklis. Implementation of efficient algorithms for globally optimal trajectories. *IEEE Trans. Automat. Control*, 43:278–283, 1998.
- [21] J. Qian, Y. Zhang, and H. Zhao. A fast sweeping method for static convex Hamilton-Jacobi equations. *J. Sci. Comput.*, 31(1-2):237–271, May 2007.
- [22] J. A. Sethian. A fast marching level set method for monotonically advancing fronts. *Proc. Natl. Acad. Sci.*, 93:1591–1595, 1996.
- [23] J. A. Sethian. *Level Set Methods*. Cambridge University Press, Cambridge, UK, 1996.
- [24] J. A. Sethian. Fast marching methods. *SIAM Review*, 41(2):199–235, 1999.
- [25] J. A. Sethian. *Level Set Methods and Fast Marching Methods*. Cambridge University Press, Cambridge, UK, 1999.
- [26] J. A. Sethian and A. Vladimirsky. Fast methods for Eikonal and related Hamilton-Jacobi equations on unstructured meshes. *Proc. Natl. Acad. Sci.*, 97(11):5699–5703, May 2000.
- [27] J. A. Sethian and A. Vladimirsky. Ordered upwind methods for static Hamilton-Jacobi equations. *Proc. Natl. Acad. Sci.*, 98(20):11069–11074, 2001.
- [28] J. A. Sethian and A. Vladimirsky. Ordered upwind methods for static Hamilton-Jacobi equa-



- tions: Theory and algorithms. *SIAM J. NUMER. ANAL.*, 41(1):323–363, 2003.
- [29] Y.-H. R. Tsai, L.-T. Cheng, S. Osher, and H.-K. Zhao. Fast sweeping algorithms for a class of Hamilton-Jacobi equations. *SIAM J. NUMER. ANAL.*, 41(2):673–694, 2003.
- [30] J. N. Tsitsiklis. Efficient algorithms for globally optimal trajectories. In *Proceedings of the 33rd Conference on Decision and Control*, pages 1368–1373, 1994.
- [31] J. N. Tsitsiklis. Efficient algorithms for globally optimal trajectories. *IEEE Trans. Automat. Control*, 40(9):1528–1538, 1995.
- [32] L. Yatziv, A. Bartesaghi, and G. Sapiro.  $O(n)$  implementation of the fast marching method. *J. Comput. Phys.*, 212(2):393–399, March 2006.
- [33] H. Zhao. A fast sweeping method for Eikonal equations. *Math. Comp.*, 74(250):603–627, 2004.