

# Particle filtering for partially observed Gaussian state space models

Christophe Andrieu

*University of Bristol, UK*

and Arnaud Doucet

*University of Cambridge, UK*

[Received October 2000. Revised March 2002]

**Summary.** Solving Bayesian estimation problems where the posterior distribution evolves over time through the accumulation of data has many applications for dynamic models. A large number of algorithms based on particle filtering methods, also known as sequential Monte Carlo algorithms, have recently been proposed to solve these problems. We propose a special particle filtering method which uses random mixtures of normal distributions to represent the posterior distributions of partially observed Gaussian state space models. This algorithm is based on a marginalization idea for improving efficiency and can lead to substantial gains over standard algorithms. It differs from previous algorithms which were only applicable to conditionally linear Gaussian state space models. Computer simulations are carried out to evaluate the performance of the proposed algorithm for dynamic tobit and probit models.

**Keywords:** Bayesian estimation; Filtering; Generalized linear time series; Importance sampling; Sequential Monte Carlo sampling; State space model

## 1. Introduction

### 1.1. Background

Many data analysis tasks involve estimating the state of a dynamic model when only partial or inaccurate observations are available (West and Harrison, 1997). Except in a few special cases, including linear Gaussian state space models, on-line state estimation is a problem that does not admit a closed form solution. As most real world models are non-linear and non-Gaussian, it is of great interest to develop efficient computational methods to solve this so-called Bayesian filtering problem numerically.

Many approximation schemes, such as the extended Kalman filter, have been proposed to surmount this problem. However, in many realistic scenarios, these approximating methods are unreliable and faults are difficult to diagnose on line. Recently there has been a surge of interest in sequential Monte Carlo (SMC) methods for non-linear or non-Gaussian time series analysis (Doucet *et al.*, 2001). These methods, initiated in Gordon *et al.* (1993), utilize a random-sample- (or particle-) based representation of the posterior probability distributions.

### 1.2. General problem

For any sequence  $l_t$ , we define

*Address for correspondence:* Arnaud Doucet, Department of Engineering, University of Cambridge, Trumpington Street, Cambridge, CB2 1PZ, UK.  
E-mail: ad2@eng.cam.ac.uk

$$l_{i:j} \triangleq (l_i, l_{i+1}, \dots, l_j).$$

In this paper, we shall concentrate on the following class of state space models. Let  $t = 1, 2, \dots$  denote discrete time: then

$$x_t = A_t x_{t-1} + B_t v_t + F_t u_t, \quad x_0 \sim \mathcal{N}(\hat{x}_0, P_0), \quad (1)$$

$$y_t = C_t x_t + D_t \varepsilon_t + G_t u_t, \quad (2)$$

$$z_t \sim p(z_t | y_t), \quad (3)$$

where  $u_t \in \mathbb{R}^{n_u}$  is an exogenous process and  $x_t \in \mathbb{R}^{n_x}$  and  $y_t \in \mathbb{R}^{n_y}$  are *unobserved* processes. The sequences

$$v_t \stackrel{\text{IID}}{\sim} \mathcal{N}(0, I_{n_v}) \in \mathbb{R}^{n_v}$$

and

$$\varepsilon_t \stackrel{\text{IID}}{\sim} \mathcal{N}(0, I_{n_\varepsilon}) \in \mathbb{R}^{n_\varepsilon}$$

are independent identically distributed (IID) Gaussian. We assume that  $P_0 > 0$ ;  $x_0$ ,  $v_t$  and  $w_t$  are mutually independent for all  $t$ , and the model parameters

$$\lambda \triangleq (\hat{x}_0, P_0, A_t, B_t, C_t, D_t, F_t, G_t; t = 1, 2, \dots)$$

are known. The processes  $(x_t)$  and  $(y_t)$  define a standard linear Gaussian state space model. We do not observe  $(y_t)$  in our case, but  $(z_t)$ . The *observations*  $(z_t)$  are conditionally independent given the processes  $(x_t)$  and  $(y_t)$  and marginally distributed according to  $p(z_t | y_t)$ ; it is assumed that  $p(z_t | y_t)$  can be evaluated pointwise up to a normalizing constant. Typically  $p(z_t | y_t)$  belongs to the exponential family. Alternatively  $z_t$  may be a censored or quantized version of  $y_t$ . This class of *partially observed Gaussian state space models* has numerous applications; many examples are discussed for instance in de Jong (1997), Manrique and Shephard (1998) and West and Harrison (1997).

We want to estimate *sequentially in time* some characteristics of the posterior distribution  $p(x_{0:t} | z_{1:t})$ . Typically, we are interested in computing  $E(x_t | z_{1:t})$  (filtering),  $E(x_{t+L} | z_{1:t})$  (prediction) and  $E(x_{t-L} | z_{1:t})$  (fixed lag smoothing), where  $L$  is a positive integer. These estimates do not in general admit analytical expressions and we must resort to numerical methods.

### 1.3. Resolution

SMC methods are, loosely speaking, a combination of importance sampling and resampling methods that allow us to propagate efficiently over time a large set of samples or particles distributed approximately according to  $p(x_{0:t} | z_{1:t})$ . We could apply standard SMC methods such as the bootstrap filter (Gordon *et al.*, 1993) to estimate  $p(x_{0:t}, y_{1:t} | z_{1:t})$  and consequently  $p(x_{0:t} | z_{1:t})$ . However, in its standard form, this algorithm does not use all the salient structure of the model. Our algorithm is based on a marginalization technique, often called the Rao–Blackwellization method (Gelfand and Smith, 1990), that improves the efficiency of the procedure. It focuses on the estimation of  $p(y_{1:t} | z_{1:t})$  rather than on the joint density  $p(x_{0:t}, y_{1:t} | z_{1:t})$ . The process  $(x_t)$  is integrated out analytically. Once  $p(y_{1:t} | z_{1:t})$  has been estimated, we can obtain estimates of  $E(x_t | z_{1:t})$ ,  $E(x_{t+L} | z_{1:t})$  and  $E(x_{t-L} | z_{1:t})$  through the Kalman filter as discussed further. In a Markov chain Monte Carlo framework, de Jong (1997) has proposed the so-called scan sampler to sample from  $p(y_{1:t} | z_{1:t})$  in a similar class of state space models; see Manrique and Shephard (1998) for some applications.

1.4. Plan

The rest of the paper is organized as follows. Section 2 shows how it is possible to restrict ourselves to estimating  $p(y_{1:t}|z_{1:t})$  instead of  $p(x_{0:t}, y_{1:t}|z_{1:t})$ , leading to an improvement in the Monte Carlo efficiency. The particle filtering algorithm is then described in detail. Section 3 demonstrates the performance of the proposed algorithm via computer simulations for dynamic tobit and probit models.

2. Rao–Blackwellized particle filtering

2.1. Marginalization

Consider the state space model defined by equations (1)–(3). We have

$$p(x_{0:t}|z_{1:t}) = \int p(x_{0:t}|y_{1:t}) p(y_{1:t}|z_{1:t}) dy_{1:t}.$$

Thus if we obtain (through an SMC method described further) an approximation of the probability distribution that is associated with the density  $p(y_{1:t}|z_{1:t})$  of the form

$$\hat{p}_N(dy_{1:t}|z_{1:t}) = \sum_{i=1}^N w_t^{(i)} \delta_{\tilde{y}_{1:t}^{(i)}}(dy_{1:t}), \quad w_t^{(i)} \geq 0, \quad \sum_{i=1}^N w_t^{(i)} = 1,$$

then  $p(x_{0:t}|z_{1:t})$  can be approximated by using

$$\hat{p}_N(x_{0:t}|z_{1:t}) = \sum_{i=1}^N w_t^{(i)} p(x_{0:t}|\tilde{y}_{1:t}^{(i)}),$$

i.e. a mixture of Gaussian densities. From such an approximation, we can estimate  $E(x_t|z_{1:t})$  and  $E(x_{t-L}|z_{1:t})$ . For example an estimate of  $E(x_t|z_{1:t})$  is given by

$$\hat{E}_N(x_t|z_{1:t}) = \int x_t \hat{p}_N(x_{0:t}|z_{1:t}) dx_{0:t} = \sum_{i=1}^N w_t^{(i)} E(x_t|\tilde{y}_{1:t}^{(i)}),$$

where  $E(x_t|\tilde{y}_{1:t}^{(i)})$  is computed through the Kalman filter associated with the linear Gaussian state space model defined by equations (1) and (2). Using the variance decomposition formula, it is clear that for any function  $h(\cdot)$

$$\text{var}\{h(x_t)|z_{1:t}\} \geq \text{var}[E\{h(x_t)|y_{1:t}, z_{1:t}\}|z_{1:t}],$$

which shows that estimating  $p(y_{1:t}|z_{1:t})$  only is more efficient.

To obtain an SMC approximation of the marginal posterior density  $p(y_{1:t}|z_{1:t})$ , we need to be able to estimate this ‘target’ density pointwise up to a normalizing constant. We have

$$p(y_{1:t}|z_{1:t}) \propto \prod_{k=1}^t p(z_k|y_k) p(y_k|y_{1:k-1}), \tag{4}$$

where

$$p(y_1|y_{1:0}) \triangleq p(y_1).$$

As  $p(z_k|y_k)$  is assumed known up to a normalizing constant, it is only necessary to estimate  $p(y_k|y_{1:k-1})$  up to a normalizing constant. This predictive density can be computed by using the Kalman filter.

The Kalman filter equations are the following. Set  $x_{0|0} = \hat{x}_0$  and  $P_{0|0} = P_0$ ; then for  $t = 1, 2, \dots$  compute

$$\left. \begin{aligned} x_{t|t-1} &= A_t x_{t-1|t-1} + F_t u_t, \\ P_{t|t-1} &= A_t P_{t-1|t-1} A_t^T + B_t B_t^T, \\ y_{t|t-1} &= C_t x_{t|t-1} + G_t u_t, \\ S_t &= C_t P_{t|t-1} C_t^T + D_t D_t^T, \\ x_{t|t} &= x_{t|t-1} + P_{t|t-1} C_t^T S_t^{-1} (y_t - y_{t|t-1}), \\ P_{t|t} &= P_{t|t-1} - P_{t|t-1} C_t^T S_t^{-1} C_t P_{t|t-1}, \end{aligned} \right\} \quad (5)$$

where

$$\begin{aligned} x_{t|t-1} &\triangleq E(x_t | y_{1:t-1}), \\ x_{t|t} &\triangleq E(x_t | y_{1:t}), \\ y_{t|t-1} &\triangleq E(y_t | y_{1:t-1}), \\ P_{t|t-1} &\triangleq \text{cov}(x_t | y_{1:t-1}), \\ P_{t|t} &\triangleq \text{cov}(x_t | y_{1:t}) \end{aligned}$$

and

$$S_t \triangleq \text{cov}(y_t | y_{1:t-1}).$$

We obtain  $p(y_k | y_{1:k-1}) = \mathcal{N}(y_k; y_{k|k-1}, S_k)$ , where  $\mathcal{N}(y_k; y_{k|k-1}, S_k)$  is a Gaussian distribution of argument  $y_k$ , mean  $y_{k|k-1}$  and covariance  $S_k$ .

## 2.2. Particle filtering

### 2.2.1. Sequential importance sampling and resampling

We describe briefly here how to apply the sequential importance sampling–resampling (SISR) method to sample approximately from  $p(y_{1:t} | z_{1:t})$ ; see Doucet *et al.* (2001) for further details.

At time  $t - 1$ , assume that we have say  $N$  particles  $\{\tilde{y}_{1:t-1}^{(i)}\}_{i=1}^N$  approximately distributed according to  $p(y_{1:t-1} | z_{1:t-1})$  and we want to obtain  $N$  particles  $\{y_{1:t}^{(i)}\}_{i=1}^N$  distributed according to  $p(y_{1:t} | z_{1:t})$ . At time  $t$ , we ‘extend’ each particle  $\tilde{y}_{1:t-1}^{(i)}$  by sampling  $\tilde{y}_t^{(i)}$  according to a conditional density  $q_t(y_t | \tilde{y}_{1:t-1}^{(i)}, z_{1:t})$ . Thus, each particle  $\tilde{y}_{1:t}^{(i)}$  is distributed according to  $p(y_{1:t-1} | z_{1:t-1}) q_t(y_t | \tilde{y}_{1:t-1}^{(i)}, z_{1:t})$ . To correct for the discrepancy between  $p(y_{1:t-1} | z_{1:t-1}) \times q_t(y_t | \tilde{y}_{1:t-1}^{(i)}, z_{1:t})$  and  $p(y_{1:t} | z_{1:t})$ , we use importance sampling so that the distribution associated with the density  $p(y_{1:t} | z_{1:t})$  is approximated by

$$\hat{p}_N(dy_{1:t} | z_{1:t}) = \frac{\sum_{i=1}^N w(\tilde{y}_{1:t}^{(i)}) \delta_{\tilde{y}_{1:t}^{(i)}}(dy_{1:t})}{\sum_{i=1}^N w(\tilde{y}_{1:t}^{(i)})} = \sum_{i=1}^N w_t^{(i)} \delta_{\tilde{y}_{1:t}^{(i)}}(dy_{1:t}), \quad (6)$$

where, using equation (4), we have for the importance weight

$$w(y_{1:t}) \propto \frac{p(y_{1:t} | z_{1:t})}{p(y_{1:t-1} | z_{1:t-1}) q_t(y_t | y_{1:t-1}, z_{1:t})} \propto \frac{p(z_t | y_t) p(y_t | y_{1:t-1})}{q_t(y_t | y_{1:t-1}, z_{1:t})}.$$

The performance of the algorithm depends on the importance density  $q_t(y_t | y_{1:t-1}, z_{1:t})$ . We can select  $p(y_t | y_{1:t-1})$  since it is a Gaussian density. In this case the associated importance weight is equal to  $w(y_{1:t}) \propto p(z_t | y_t)$ . Note that the ‘optimal’ importance density, i.e. the density minimizing the conditional variance of the weight conditional on  $y_{1:t-1}$  (Doucet *et al.*, 2000), is

$$p(y_t | y_{1:t-1}, z_{1:t}) \propto p(z_t | y_t) p(y_t | y_{1:t-1}),$$

and the associated importance weight is

$$w(y_{1:t}) \propto p(z_t | y_{1:t-1}) = \int p(z_t | y_t) p(y_t | y_{1:t-1}) dy_t. \tag{7}$$

It might be possible to compute this weight or not, depending on  $p(z_t | y_t)$ .

Finally, we obtain  $N$  particles  $\{y_{1:t}^{(i)}\}_{i=1}^N$  approximately distributed according to  $p(y_{1:t} | z_{1:t})$  by resampling from the weighted empirical distribution given in equation (6). Several resampling procedures are available in the literature. We adopt here the stratified sampling scheme described in Kitagawa (1996).

Alternative SMC methods can be applied to estimate  $p(y_{1:t} | z_{1:t})$ . In particular, the auxiliary particle filtering (APF) technique of Pitt and Shephard (1999) could be used. The idea behind APF is to extend existing particles  $\tilde{y}_{1:t-1}^{(i)}$  that are the most promising, in the sense that the predictive likelihood  $p(z_t | \tilde{y}_{1:t-1}^{(i)})$  is large. When  $p(z_t | \tilde{y}_{1:t-1}^{(i)})$  cannot be computed analytically, APF proposes an analytical approximation. In this case, APF and SISR differ significantly. However, when  $p(z_t | \tilde{y}_{1:t-1}^{(i)})$  can be computed analytically, then APF uses the optimal importance density. This is referred to as ‘perfect adaptation’ (Pitt and Shephard, 1999). In this particular case, APF and SISR are essentially similar except that APF reverses the order of the sampling and resampling steps; this is possible as the importance weight is independent of  $y_t$  and this is obviously more efficient.

### 2.2.2. Algorithm

We limit our presentation to standard choices of importance densities where  $q_t(y_t | y_{1:t-1}, z_{1:t})$  depends on  $(y_{1:t-1}, z_{1:t})$  only via  $z_t$  and the set of low-dimensional sufficient statistics  $x_{t|t-1}$  and  $P_{t|t-1}$ . We shall write

$$q_t(y_t | x_{t|t-1}, P_{t|t-1}, z_t) \triangleq q_t(y_t | y_{1:t-1}, z_{1:t}).$$

This class of densities includes

$$p(y_t | x_{t|t-1}, P_{t|t-1}) \triangleq p(y_t | y_{1:t-1}) = \mathcal{N}(y_t; y_{t|t-1}, S_t),$$

where  $y_{t|t-1}$  and  $S_t$  are deterministic functions of  $x_{t|t-1}$  and  $P_{t|t-1}$ . As one typically focuses on features of the marginal  $p(y_t | z_{1:t})$ , it is only necessary to store in memory  $\{y_t^{(i)}, x_{t|t-1}^{(i)}\}_{i=1}^N$  and  $P_{t|t-1}$  instead of  $\{y_{1:t}^{(i)}\}_{i=1}^N$ . Contrary to algorithms presented in Chen and Liu (2000) and Doucet *et al.* (2000), we point out that we do not have to compute  $N$  ‘full’ Kalman filter recursions as most of the calculations need to be done only once. More precisely, we note that  $P_{t|t-1}^{(i)} = P_{t|t-1}$  and  $S_{t|t}^{(i)} = S_{t|t}$  for any  $i \in \{1, \dots, N\}$ .

Given  $N$  particles  $\{y_{t-1}^{(i)}\}_{i=1}^N$  at time  $t-1$  distributed approximately according to  $p(y_{t-1} | z_{1:t-1})$  and the associated sufficient statistics  $\{x_{t-1|t-1}^{(i)}\}_{i=1}^N$  and  $P_{t-1|t-1}$ , the particle filter proceeds as follows at time  $t$ . In the *sequential importance sampling step*:

- (a) for  $i = 1, \dots, N$ , set

$$\tilde{x}_{t|t-1}^{(i)} \triangleq x_{t|t-1}^{(i)}$$

and sample

$$\tilde{y}_t^{(i)} \sim q_t(y_t | \tilde{x}_{t|t-1}^{(i)}, P_{t|t-1}, z_t);$$

(b) for  $i = 1, \dots, N$ , evaluate and normalize the importance weights

$$w_t^{(i)} \propto \frac{p(z_t | \tilde{y}_t^{(i)}) p(\tilde{y}_t^{(i)} | \tilde{x}_{t|t-1}^{(i)}, P_{t|t-1})}{q_t(\tilde{y}_t^{(i)} | \tilde{x}_{t|t-1}^{(i)}, P_{t|t-1}, z_t)}, \quad \sum_{i=1}^N w_t^{(i)} = 1. \tag{8}$$

In the *resampling step*: multiply or discard particles  $\{\tilde{y}_t^{(i)}, \tilde{x}_{t|t-1}^{(i)}\}_{i=1}^N$  with respect to high or low importance weights  $w_t^{(i)}$  to obtain  $N$  particles  $\{y_t^{(i)}, x_{t|t-1}^{(i)}\}_{i=1}^N$ .

In the *updating step*:

- (a) compute  $P_{t+1|t}$  given  $P_{t|t-1}$  using one step of the Kalman recursion (5);
- (b) for  $i = 1, \dots, N$ , use one step of the Kalman recursion (5) to compute  $x_{t+1|t}^{(i)}$  given  $y_t^{(i)}, x_{t|t-1}^{(i)}$  and  $P_{t|t-1}$ .

The computational complexity of this algorithm at each time step is  $\mathcal{O}(N)$ . If the (unnormalized) importance weights given by equation (8) are upper bounded, then asymptotic convergence ( $N \rightarrow \infty$ ) of Monte Carlo estimates towards their true values can be ensured (Crisan, 2001).

### 2.3. Extensions

There are many potential extensions to both the model and the algorithm.

#### 2.3.1. Model

For the linear Gaussian model (1)–(2), we can readily consider the case where  $v_t$  and  $w_t$  are correlated and/or add a non-linear term  $\varphi(y_{1:t-1})$  to the right-hand side of equation (2). It is also possible to apply the marginalization method presented above to integrate out analytically ( $x_t$ ) when the model (1)–(2) is not linear Gaussian but conditionally linear Gaussian as described in Shephard (1994); this extension allows us to consider finite or continuous mixtures of Gaussian distributions.

Another interesting extension consists of partially observed hidden Markov models: ( $x_t$ ) is modelled as a finite state space Markov chain and

$$p(y_{1:t}, z_{1:t} | x_{1:t}) = \prod_{k=1}^t p(y_k | x_k, y_{1:k-1}) p(z_k | y_k).$$

We can integrate out ( $x_t$ ) and compute  $p(y_k | y_{1:k-1})$  by using the hidden Markov model filter instead of the Kalman filter as part of the method developed in the previous section.

#### 2.3.2. Algorithm

When the distribution of the importance weights  $w_t^{(i)}$  is skewed, the particles  $\{\tilde{y}_{1:t}^{(i)}\}_{i=1}^N$  having high importance weights are selected many times; this results in a ‘depletion’ of samples as numerous particles  $y_{1:t}^{(i)}$  and  $y_{1:t}^{(j)}$  are in fact equal for  $i \neq j$ . To perform sample ‘regeneration’, it is possible to use a recently proposed approach based on Markov chain Monte Carlo steps (Gilks and Berzuini, 2001). It consists of applying to each particle  $y_{1:t}^{(i)}$  a (possibly non-ergodic) transition kernel  $K_t(y_{1:t} | y_{1:t}^{(i)})$  of invariant density  $p(y_{1:t} | z_{1:t})$ . There are an infinite number of possible choices for this kernel. One possibility consists of updating at time  $t$  the values  $y_{t-M+1:t}$  ( $M > 0$ ) by using the efficient scan sampler (de Jong, 1997), whose computational complexity is  $\mathcal{O}(M)$ . Though this step is not necessary to ensure theoretical convergence of the algorithm, it can improve results.

### 3. Simulations

#### 3.1. Dynamic tobit model

Let us consider the following tobit model (Manrique and Shephard, 1998):

$$\begin{aligned} x_{t+1} &= \phi x_t + \sigma_v v_{t+1}, & x_0 &\sim \mathcal{N}\{0, \sigma_v^2/(1 - \phi^2)\}, & v_t &\stackrel{\text{IID}}{\sim} \mathcal{N}(0, 1), \\ y_t &= x_t + \sigma_\varepsilon \varepsilon_t, & \varepsilon_t &\stackrel{\text{IID}}{\sim} \mathcal{N}(0, 1), \\ z_t &= \max(y_t, 0). \end{aligned}$$

It is clear that this model is of the form (1)–(3). We chose as importance density the ‘optimal’ density  $p(y_t | y_{1:t-1}, z_t)$ . (We could not use  $p(y_t | y_{1:t-1})$  for the importance density when  $z_t = 0$ . In this case, the importance weight does not exist.) If  $z_t > 0$ , then  $y_t = z_t$  and, if  $z_t = 0$ , then

$$p(y_t | y_{1:t-1}, z_t = 0) \propto p(y_t | y_{1:t-1}) \mathbb{1}_{(-\infty, 0)}(y_t).$$

For the importance weight, we obtain using equation (7)

$$w(y_{1:t}) \propto \begin{cases} \Phi(-y_{t|t-1}/\sqrt{S_t}) & \text{if } z_t = 0, \\ \mathcal{N}(z_t; y_{t|t-1}, S_t) & \text{if } z_t > 0, \end{cases}$$

where  $\Phi(\cdot)$  is the cumulative function of the standard normal distribution.

We simulated  $T = 200$  observations with the known hyperparameters  $\phi = 0.99$ ,  $\sigma_v^2 = 0.05$  and  $\sigma_w^2 = 0.30$ . For various numbers of particles  $N$ , we generated  $K = 100$  different realizations of our Rao–Blackwellized (RB) filter and of the standard algorithm that estimates  $p(x_{0:t}, y_{1:t} | z_{1:t})$  using the importance density  $p(x_t, y_t | x_{t-1}, y_{t-1}, z_t)$  as  $p(x_t, y_t | x_{t-1}, y_{t-1})$  cannot be used. Our comparison is in terms of the mean and variance of the square error SE computed as follows:

$$\begin{aligned} \text{SE}(l) &= \sum_{t=1}^T \{x_t - \hat{E}_{N,l}(x_t | z_{1:t})\}^2, \\ m(\text{SE}) &= \frac{1}{K} \sum_{l=1}^K \text{SE}(l), \\ \sigma^2(\text{SE}) &= \frac{1}{K} \sum_{l=1}^K \{\text{SE}(l) - m(\text{SE})\}^2, \end{aligned}$$

where  $\hat{E}_{N,l}(x_t | z_{1:t})$  is computed using the  $l$ th realization of the particle filter. We present in Tables 1 and 2 the performance of the standard and RB filters. For a fixed number of particles, the RB filter is not more computationally intensive than the standard filter and it performs significantly better.

#### 3.2. Dynamic probit model

We analyse here a non-stationary binary time series and more specifically the (aggregated) Tokyo rainfall data set (Knorr-Held, 1999). It consists of  $T = 366$  observations with  $z_t = 1$  indicating that it rained on the  $t$ th day of the year and  $z_t = 0$  otherwise. We model  $z_t$  by using a dynamic probit model, i.e.

$$\Pr(z_t = 1 | \alpha_t) = \Phi(\alpha_t),$$

where  $(\alpha_t)$  is modelled by using a second-order random walk

$$\alpha_t = 2\alpha_{t-1} - \alpha_{t-2} + \sigma_v v_t, \quad v_t \stackrel{\text{IID}}{\sim} \mathcal{N}(0, 1). \tag{9}$$

**Table 1.**  $m(\text{SE})$  for the standard filter and RB filter

Algorithm	Results for the following values of $N$ :							
	100	250	500	1000	2500	5000	10000	25000
Standard filter	33.70	33.64	33.90	33.41	33.45	33.55	33.54	33.52
RB filter	33.52	33.49	33.51	33.50	33.50	33.49	33.51	33.50

Let us introduce an artificial latent process  $(y_t)$  such that

$$y_t = \alpha_t + \varepsilon_t, \quad \varepsilon_t \stackrel{\text{IID}}{\sim} \mathcal{N}(0, 1), \tag{10}$$

and define

$$z_t = \mathbb{1}_{[0, \infty)}(y_t). \tag{11}$$

It is easy to check that we have

$$\Pr(z_t = 1 | \alpha_t) = \Pr(y_t > 0 | \alpha_t) = \Pr(\varepsilon_t > -\alpha_t) = \Phi(\alpha_t).$$

We can easily rewrite equations (9)–(11) in a state space model of the form (1)–(3) by defining

$$x_t \triangleq (\alpha_t, \alpha_{t-1}).$$

In this case, as the introduction of  $(y_t)$  is artificial, it is necessary to compare our procedure with a particle filtering method applied to the estimation of  $p(x_{0:t} | z_{1:t})$  and not with  $p(x_{0:t}, y_{1:t} | z_{1:t})$ . The motivation for introducing  $(y_t)$  comes from the fact that it is possible to use the optimal density as importance density

$$p(y_t | y_{1:t-1}, z_t) \propto \begin{cases} p(y_t | y_{1:t-1}) \mathbb{1}_{[0, \infty)}(y_t) & \text{if } z_t = 1, \\ p(y_t | y_{1:t-1}) \mathbb{1}_{(-\infty, 0)}(y_t) & \text{if } z_t = 0, \end{cases}$$


which is a truncated Gaussian distribution, and, using equation (7), we obtain

$$w(y_{1:t}) \propto p(z_t | y_{1:t-1}) = \left\{ 1 - \Phi\left(-\frac{y_{t|t-1}}{\sqrt{S_t}}\right) \right\}^{z_t} \Phi\left(-\frac{y_{t|t-1}}{\sqrt{S_t}}\right)^{1-z_t}.$$

If we consider  $p(x_{0:t} | z_{1:t})$ , the optimal density cannot be used since the associated importance weight  $p(z_t | x_{t-1})$  does not admit an analytical expression. Note that the introduction of  $(y_t)$

**Table 2.**  $10 \sigma(\text{SE})$  for the standard filter and RB filter

Algorithm	Results for the following values of $N$ :							
	100	250	500	1000	2500	5000	10000	25000
Standard filter	3.76	2.19	1.61	1.20	0.83	0.62	0.37	0.26
RB filter	2.50	1.30	1.21	0.99	0.51	0.38	0.29	0.14



**Fig. 1.** (a) Binary observations ( $z_t$ ) and (b)  $E\{\Phi(\alpha_t)|y_{1:t}\}$  (—) and  $E\{\Phi(\alpha_t)|y_{1:t}\} \pm \sqrt{\text{var}\{\Phi(\alpha_t)|y_{1:t}\}}$  (---)

has already been used to develop efficient Markov chain Monte Carlo samplers (Albert and Chib, 1993).

The hyperparameter  $\sigma_v^2$  was set to 0.01. In Fig. 1, we display the observations  $z_t$ ,  $E\{\Phi(\alpha_t)|y_{1:t}\}$  and  $E\{\Phi(\alpha_t)|y_{1:t}\} \pm \sqrt{\text{var}\{\Phi(\alpha_t)|y_{1:t}\}}$ ; the estimates are obtained by using  $N = 1000$  particles. To obtain similar results, the bootstrap filter (Gordon *et al.*, 1993) requires as many as  $N = 5000$  particles.

#### 4. Conclusion

This paper has proposed a method for recursive state estimation of partially observed Gaussian models. Our algorithm is an SMC method based on marginalization. This marginalization is performed through Kalman filtering methods. Our simulations show that our approach can significantly outperform standard SMC methods.

Throughout the paper, the model parameters  $\lambda$  are assumed known. It is possible, however, to perform batch and recursive estimation of these parameters, combining the particle filtering method developed here and recent methods proposed in Liu and West (2001).

#### Acknowledgements

We are grateful to Leonhard Knorr-Held for providing the Tokyo rainfall data set and to Sanjeev Arulampalam, Nando de Freitas, Arthur Gretton and Elena Punskeya for valuable comments.

#### References

- Albert, J. and Chib, S. (1993) Bayesian analysis of binary and polychotomous response data. *J. Am. Statist. Ass.*, **88**, 669–679.  
 Chen, R. and Liu, J. S. (2000) Mixture Kalman filters. *J. R. Statist. Soc. B*, **62**, 493–508.

- Crisan, D. (2001) Particle filters—a theoretical perspective. In *Sequential Monte Carlo Methods in Practice* (eds A. Doucet, J. F. G. de Freitas and N. J. Gordon), pp. 17–41. New York: Springer.
- Doucet, A., de Freitas, J. F. G. and Gordon, N. J. (eds) (2001) *Sequential Monte Carlo Methods in Practice*. New York: Springer.
- Doucet, A., Godsill, S. J. and Andrieu, C. (2000) On sequential Monte Carlo sampling methods for Bayesian filtering. *Statist. Comput.*, **10**, 197–208.
- Gelfand, A. and Smith, A. F. M. (1990) Sampling based approaches to calculating marginal densities. *J. Am. Statist. Ass.*, **85**, 398–409.
- Gilks, W. R. and Berzuini, C. (2001) Following a moving target—Monte Carlo inference for dynamic Bayesian models. *J. R. Statist. Soc. B*, **63**, 127–146.
- Gordon, N. J., Salmond, D. J. and Smith, A. F. M. (1993) Novel approach to nonlinear/non-Gaussian Bayesian state estimation. *IEE Proc. F*, **140**, 107–113.
- de Jong, P. (1997) The scan sampler. *Biometrika*, **84**, 929–937.
- Kitagawa, G. (1996) Monte Carlo filter and smoother for non-Gaussian nonlinear state space models. *J. Comput. Graph. Statist.*, **5**, 1–25.
- Knorr-Held, L. (1999) Conditional prior proposals in dynamic models. *Scand. J. Statist.*, **26**, 129–144.
- Liu, J. and West, M. (2001) Combined parameter and state estimation in simulation-based filtering. In *Sequential Monte Carlo Methods in Practice* (eds A. Doucet, J. F. G. de Freitas and N. J. Gordon), pp. 197–223. New York: Springer.
- Manrique, A. and Shephard, N. (1998) Simulation-based likelihood inference for limited dependent processes. *Econometr. J.*, **1**, 174–202.
- Pitt, M. K. and Shephard, N. (1999) Filtering via simulation: auxiliary particle filter. *J. Am. Statist. Ass.*, **94**, 590–599.
- Shephard, N. (1994) Partial non-Gaussian times series models. *Biometrika*, **81**, 115–131.
- West, M. and Harrison, P. J. (1997) *Bayesian Forecasting and Dynamic Models*, 2nd edn. New York: Springer.