

# Reasoning Under Uncertainty: Bnet Inference

(Variable elimination)

Computer Science cpsc322, Lecture 29

*(Textbook Chpt 6.4)*




Nov, 16, 2012

# Lecture Overview



- **Recap Learning Goals previous lecture**
- Bnets Inference
  - Intro
  - Factors
  - Variable elimination Intro

# Learning Goals for Wed's class

You can:


- In a Belief Net, determine whether one variable is independent of another variable, given a set of observations.
  - Define and use **Noisy-OR** distributions. Explain assumptions and benefit.
  - Implement and use a **naïve Bayesian classifier**. Explain assumptions and benefit.
- 
- A hand-drawn blue bracket on the right side of the slide groups the last two bullet points. There are also two hand-drawn blue arrows: one pointing to the 'Noisy-OR' text in the second bullet point, and another pointing to the 'naïve Bayesian classifier' text in the third bullet point.

### 3 Configuration blocking dependency (belief propagation)



# Bnets: Compact Representations

n Boolean variables, k max. number of parents



Only one parent with  $h$  possible values




# Lecture Overview

- Recap Learning Goals previous lecture
- **Bnets Inference**
  - Intro
  - Factors
  - Variable elimination Algo

# Bnet Inference

- **Our goal:** compute probabilities of variables in a belief network

What is the posterior distribution over one or more variables, conditioned on one or more observed variables?



*examples*

$$\begin{aligned} &\rightarrow P(\text{Alarm} \mid \text{Smoke} = f) \\ &P(\text{Fire} \mid \text{Smoke} = t, \text{Leaving} = f) \end{aligned}$$



# Bnet Inference: General

- Suppose the variables of the belief network are  $X_1, \dots, X_n$ .
- $Z$  is the query variable
- $Y_1=v_1, \dots, Y_j=v_j$  are the observed variables (with their values)
- $Z_1, \dots, Z_k$  are the remaining variables
- What we want to compute:  $P(Z \mid Y_1 = v_1, \dots, Y_j = v_j)$   $\swarrow$



*Example:*

$$P(L \mid S = t, R = f) \swarrow$$

$$Z \leftrightarrow L$$

$$Y_1 Y_2 \leftrightarrow S, R$$

$$Z_1 Z_2 Z_3 \leftrightarrow T, F, A$$



# What do we need to compute?

Remember conditioning and marginalization...

$$P(L | S=t, R=f) = \frac{P(L, S=t, R=f) \leftarrow \textcircled{1}}{P(S=t, R=f) \textcircled{2}}$$

| L | S | R | $P(L, S=t, R=f)$ |
|---|---|---|------------------|
| t | t | f | .3               |
| f | t | f | .2               |

*Do they have to sum up to one?*

no

yes

no

$$\textcircled{2} = .5$$

| L | S | R | $P(L   S=t, R=f)$ |
|---|---|---|-------------------|
| t | t | f | .6                |
| f | t | f | .4                |

# In general.....

$$\underbrace{P(Z | Y_1 = v_1, \dots, Y_j = v_j)} = \frac{\boxed{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}}{\boxed{P(Y_1 = v_1, \dots, Y_j = v_j)}} \Rightarrow \frac{\boxed{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}}{\sum_Z \boxed{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}}$$

①  
②

- We only need to **compute the** *numerator* and then **normalize**
- This can be framed in terms of **operations between factors** (that satisfy the semantics of probability)

# Lecture Overview

- Recap Bnets
- Bnets Inference
  - Intro
  - **Factors**
  - Variable elimination Algo

# Factors

- A **factor** is a representation of a function from a tuple of random variables into a number.  $[0, 1]$

- We will write factor  $f$  on variables  $X_1, \dots, X_j$  as  $f(X_1, \dots, X_j)$

- A factor denotes one or more (possibly partial) distributions over the given tuple of variables

- e.g.,  $P(X_1, X_2)$  is a factor  $f(X_1, X_2)$  *Distribution*

- e.g.,  $P(X_1, X_2, X_3 = v_3)$  is a factor  $f(X_1, X_2)_{X_3 = v_3}$  *Partial distribution*

- e.g.,  $P(Z | X, Y)$  is a factor  $f(Z, X, Y)$  *Set of Distributions*

- e.g.,  $P(X_1, X_3 = v_3 | X_2)$  is a factor  $f(X_1, X_2)_{X_3 = v_3}$  *Set of partial Distributions*

$$P(Z | X, Y)$$

| X | Y | Z | val |
|---|---|---|-----|
| t | t | t | 0.1 |
| t | t | f | 0.9 |
| t | f | t | 0.2 |
| t | f | f | 0.8 |
| f | t | t | 0.4 |
| f | t | f | 0.6 |
| f | f | t | 0.3 |
| f | f | f | 0.7 |

# Factors

- A **factor** is a representation of a function from a tuple of random variables into a number.  $[0, 1]$
- We will write factor  $f$  on variables  $X_1, \dots, X_j$  as  $f(X_1, \dots, X_j)$
- A factor denotes one or more (possibly partial) distributions over the given tuple of variables

$P(Z|X,Y)$

- e.g.,  $P(X_1, X_2)$  is a factor  $f(X_1, X_2)$  *Distribution*
- e.g.,  $P(X_1, X_2, X_3 = v_3)$  is a factor  $f(X_1, X_2)_{X_3 = v_3}$  *Partial distribution*
- e.g.,  $P(X | Z, Y)$  is a factor  $f(X, Z, Y)$  *Set of Distributions*
- e.g.,  $P(X_1, X_3 = v_3 | X_2)$  is a factor  $f(X_1, X_2)_{X_3 = v_3}$  *Set of partial Distributions*

| $P(X,Y,Z)$ |   | $P(Y Z,X)$ |     |
|------------|---|------------|-----|
| X          | Y | Z          | val |
| t          | t | t          | 0.1 |
| t          | t | f          | 0.9 |
| t          | f | t          | 0.2 |
| t          | f | f          | 0.8 |
| f          | t | t          | 0.4 |
| f          | t | f          | 0.6 |
| f          | f | t          | 0.3 |
| f          | f | f          | 0.7 |

# Factors

- A **factor** is a representation of a function from a tuple of random variables into a number.  $[0, 1]$

- We will write factor  $f$  on variables  $X_1, \dots, X_j$  as  $f(X_1, \dots, X_j)$

- A factor denotes one or more (possibly partial) distributions over the given tuple of variables

$$P(Z|X,Y)$$

- e.g.,  $P(X_1, X_2)$  is a factor  $f(X_1, X_2)$  *Distribution*

- e.g.,  $P(X_1, X_2, X_3 = v_3)$  is a factor *Partial distribution*  
 $f(X_1, X_2)_{X_3 = v_3}$

- e.g.,  $P(X | Z, Y)$  is a factor *Set of Distributions*  
 $f(X, Z, Y)$

- e.g.,  $P(X_1, X_3 = v_3 | X_2)$  is a factor *Set of partial Distributions*  
 $f(X_1, X_2)_{X_3 = v_3}$


| X | Y | Z | val |
|---|---|---|-----|
| t | t | t | 0.1 |
| t | t | f | 0.9 |
| t | f | t | 0.2 |
| t | f | f | 0.8 |
| f | t | t | 0.4 |
| f | t | f | 0.6 |
| f | f | t | 0.3 |
| f | f | f | 0.7 |

# Manipulating Factors:

We can make new factors out of an existing factor

- Our first operation: we can assign some or all of the variables of a factor.

$f(X,Y,Z):$



| X | Y | Z | val |
|---|---|---|-----|
| t | t | t | 0.1 |
| t | t | f | 0.9 |
| t | f | t | 0.2 |
| t | f | f | 0.8 |
| f | t | t | 0.4 |
| f | t | f | 0.6 |
| f | f | t | 0.3 |
| f | f | f | 0.7 |

*What is the result of  
assigning  $X=t$  ?*

$f(X=t, Y, Z)$

$f(X, Y, Z)_{X=t}$

# More examples of assignment

$r(X,Y,Z):$

| X | Y | Z | val |
|---|---|---|-----|
| t | t | t | 0.1 |
| t | t | f | 0.9 |
| t | f | t | 0.2 |
| t | f | f | 0.8 |
| f | t | t | 0.4 |
| f | t | f | 0.6 |
| f | f | t | 0.3 |
| f | f | f | 0.7 |

$r(X=t,Y,Z):$



| Y            | Z            | val            |
|--------------|--------------|----------------|
| <del>t</del> | <del>t</del> | <del>0.1</del> |
| t            | f            | 0.9            |
| <del>f</del> | <del>t</del> | <del>0.2</del> |
| f            | f            | 0.8            |

$r(X=t,Y,Z=f):$

| Y            | val           |
|--------------|---------------|
| <del>t</del> | <del>.1</del> |
| f            | .8            |

$r(X=t,Y=f,Z=f):$

| val |
|-----|
| .8  |



# Summing out a variable example

Our second operation: we can *sum out* a variable, say  $X_1$  with domain  $\{v_1, \dots, v_k\}$ , from factor  $f(X_1, \dots, X_j)$ , resulting in a factor on  $X_2, \dots, X_j$  defined by:

$f_3(A,B,C):$

| B | A | C | val  |
|---|---|---|------|
| t | t | t | 0.03 |
| t | t | f | 0.07 |
| f | t | t | 0.54 |
| f | t | f | 0.36 |
| t | f | t | 0.06 |
| t | f | f | 0.14 |
| f | f | t | 0.48 |
| f | f | f | 0.32 |

$\sum_B f_3(A,C):$

| A | C | val |
|---|---|-----|
| t | t | .57 |
| t | f | .43 |
| f | t |     |
| f | f |     |

$$\left( \sum_{X_1} f \right) (X_2, \dots, X_j) = f(X_1 = v_1, X_2, \dots, X_j) + \dots + f(X_1 = v_k, X_2, \dots, X_j)$$

# Multiplying factors

- Our third operation: factors can be *multiplied* together.

$f_1(A,B)$ :

| A | B | Val |
|---|---|-----|
| t | t | 0.1 |
| t | f | 0.9 |
| f | t | 0.2 |
| f | f | 0.8 |

$f_2(B,C)$ :

| B | C | Val |
|---|---|-----|
| t | t | 0.3 |
| t | f | 0.7 |
| f | t | 0.6 |
| f | f | 0.4 |

$f_1(A,B) \times f_2(B,C)$ :

| A | B | C | val  |
|---|---|---|------|
| t | t | t | .03  |
| t | t | f | .07  |
| t | f | t | .054 |
| t | f | f |      |
| f | t | t |      |
| f | t | f |      |
| f | f | t |      |
| f | f | f |      |

# Multiplying factors: Formal

- The **product** of factor  $f_1(A, B)$  and  $f_2(B, C)$ , where  $B$  is the variable in common, is the factor  $(f_1 \times f_2)(A, B, C)$  defined by:

$$f_1(A, B) f_2(B, C) = (f_1 \times f_2)(A, B, C)$$

$\begin{matrix} t & f & f & & \\ & & & t & f \\ & & & A & B & & B & C \end{matrix}$

**Note1:** it's defined on all  $A, B, C$  triples, obtained by multiplying together the appropriate pair of entries from  $f_1$  and  $f_2$ .

**Note2:**  $A, \textcircled{B}, C$  can be sets of variables

# Factors Summary

- A factor is a representation of a function from a tuple of random variables into a number.
  - $f(X_1, \dots, X_j)$ .
- We have defined three operations on factors:
  1. Assigning one or more variables
    - $f(X_1=v_1, X_2, \dots, X_j)$  is a factor on  $X_2, \dots, X_j$ , also written as  $f(X_1, \dots, X_j)_{X_1=v_1}$
  2. Summing out variables
    - $(\sum_{X_1} f)(X_2, \dots, X_j) = f(X_1=v_1, X_2, \dots, X_j) + \dots + f(X_1=v_k, X_2, \dots, X_j)$
  3. Multiplying factors
    - $f_1(A, B) f_2(B, C) = (f_1 \times f_2)(A, B, C)$

# Lecture Overview

- Recap Bnets
- Bnets Inference
  - Intro
  - Factors
  - Intro Variable elimination Algo

# Variable Elimination Intro

- Suppose the variables of the belief network are  $X_1, \dots, X_n$ .
- $Z$  is the query variable
- $Y_1=v_1, \dots, Y_j=v_j$  are the observed variables (with their values)
- $Z_1, \dots, Z_k$  are the remaining variables
- What we want to compute:  $P(Z \mid Y_1 = v_1, \dots, Y_j = v_j)$
- We showed before that what we actually need to compute is

$$P(Z, Y_1 = v_1, \dots, Y_j = v_j)$$

This can be computed in terms of **operations between factors** (that satisfy the semantics of probability)

# Variable Elimination Intro

- If we express the joint as a factor,

$$f(Z, \overbrace{Y_1, \dots, Y_j}^{\text{observed}}, \underbrace{Z_1, \dots, Z_k}_{\text{sum out}})$$

assign

- We can compute  $P(Z, Y_1=v_1, \dots, Y_j=v_j)$  by ??

- assigning  $Y_1=v_1, \dots, Y_j=v_j$

- and summing out the variables  $Z_1, \dots, Z_k$

$$P(Z, Y_1=v_1, \dots, Y_j=v_j) = \sum_{Z_k} \cdots \sum_{Z_1} f(Z, Y_1, \dots, Y_j, Z_1, \dots, Z_k)_{Y_1=v_1, \dots, Y_j=v_j}$$

*Are we done?*

no

this is the joint Too BIG!

# Learning Goals for today's class

You can:

- Define **factors**. Derive new factors from existing factors. Apply operations to factors, including assigning, summing out and multiplying factors.
- (*Minimally*) Carry out **variable elimination** by using factor representation and using the factor operations. Use techniques to simplify variable elimination.



# Next Class

## Variable Elimination

- The algorithm
- An example

## Course Elements

- Assignment 3 is due on Monday!
- Assignment 4 will be available on Wednesday and due on Nov the 28th (last class).