


# Course Updates

- Midterm is today @ 5pm



# Lighting and Reflections

---

Computer Graphics, CSCD18

Fall 2007

Instructor: Leonid Sigal



# Introduction

- So far we have only considered in our rendering pipeline
  - Geometry of the scene
  - Camera modeling
- We did not consider
  - Lighting
  - Shading of polygons (must account for properties of the surface)
  - Shadows

# Introduction

- In general, to reason about shading, lighting and shadows
  - We must consider every light ray that hits every surface in the scene
    - Rays that come directly from the light source
    - Rays that are reflected from different object surfaces
  - This is impractical for most scenes
  - We need to make simplifications


# Lighting Basics

## ■ Light sources

- Point sources (e.g. sun)
  - Light is reflected in all directions from the small light source far away
- Extended light sources (e.g. day lights)
  - Light is reflected in all directions but from many points
- Directional lighting
- Secondary lighting
  - Light reflected from other objects

# Reflection Basics

- Reflectance (different objects reflect light in different ways)
  - Diffuse surfaces (e.g. egg)
    - Appear the same from all directions
  - Secular surfaces (e.g. mirrors)
    - Reflected light is a function of the viewing direction
  - Transmission (e.g. skin, glass, water)
    - Light can penetrate the surface



# Diffuse Reflection


- **Idea:** reflected light is the same in all directions
- **Assumptions:** point light source
- Simplest model:  $L_d(\bar{p}) = r_d I \max(0, \vec{s} \cdot \vec{n})$

$I$  - intensity of the light source

$r_d$  - fraction of the light being reflected

$\vec{s}$  - direction of the light source

$\vec{n}$  - surface normal



# Diffuse Reflection


- **Idea:** reflected light is the same in all directions
- **Assumptions:** point light source
- Simplest model:  $L_d(\bar{p}) = r_d I \max(0, \vec{s} \cdot \vec{n})$  Why?

$I$  - intensity of the light source

$r_d$  - fraction of the light being reflected

$\vec{s}$  - direction of the light source (normalized)

$\vec{n}$  - surface normal (normalized)





# Diffuse Reflection


- **Idea:** reflected light is the same in all directions
- **Assumptions:** point light source
- Simplest model:  $L_d(\bar{p}) = r_d I \max(0, \vec{s} \cdot \vec{n})$

$I$  - intensity of the light source

$r_d$  - fraction of the light being reflected

$\vec{s}$  - direction of the light source (normalized)

$\vec{n}$  - surface normal (normalized)



# Diffuse Reflection with Multiple Lights

- **Idea:** reflected light is the same in all directions
- **Assumptions:** point light source


- Light is additive, so  $L_d(\bar{p}) = \sum_j r_d I_j \max(0, \vec{s}_j \cdot \vec{n})$

$I_j$  - intensity of the light source  $j$

$r_d$  - fraction of the light being reflected

$\vec{s}_j$  - direction of the light source  $j$  (normalized)

$\vec{n}$  - surface normal (normalized)




# What about color?

- **Idea:** reflected light is the same in all directions
- **Assumptions:** point light source
- We can specify reflectance and light intensity in terms of color components

$$L_{d,R}(\bar{p}) = r_{d,R} I_{j,R} \max(0, \vec{s} \cdot \vec{n})$$

$$L_{d,G}(\bar{p}) = r_{d,G} I_{j,G} \max(0, \vec{s} \cdot \vec{n})$$

$$L_{d,B}(\bar{p}) = r_{d,B} I_{j,B} \max(0, \vec{s} \cdot \vec{n})$$



# Ambient Illumination

- Diffuse reflection with point light source produce strong shadows
- Surface patches that point away from the light source,  $\vec{s} \cdot \vec{n} < 0$ , end up being black. This looks unnatural. **Why?**
- Solutions
  - Have many light sources to approximate an extended light source
  - Use ambient reflectance
    - Approximates the average amount of light in the scene

# Ambient Illumination

- Simple ambient reflectance

$$L_a(\bar{p}) = r_a I_a$$

$I_a$  - amount of ambient illumination

$r_a$  - ambient reflection coefficient (how much light is reflected)  
(often people set  $r_a = r_d$ )

- Color ambient reflectance

$$L_{a,R}(\bar{p}) = r_{a,R} I_{a,R}$$


$$L_{a,G}(\bar{p}) = r_{a,G} I_{a,G}$$

$$L_{a,B}(\bar{p}) = r_{a,B} I_{a,B}$$

# Diffuse Reflectance Model




Increasing ratio of ambient to diffuse reflection: 



# Specular Reflectance

- **Idealization:** a mirror
- Models plastics, metals, and polished surfaces
- **Property:** Angle of reflection equal to the angle of incident with respect to the normal


$\vec{r}$  - is unit vector corresponding to emitting direction  
(it is determined by the normal and the light source)



# Specular Reflectance

- **Idealization:** a mirror
- Models plastics, metals, and polished surfaces
- **Property:** Angle of reflection equal to the angle of incident with respect to the normal

$$\vec{r} = 2(\vec{n} \cdot \vec{s})\vec{n} - \vec{s}$$



In practice most specular surfaces reflect light close to this direction



# Specular Reflectance

## ■ Common specular model

$$L_s(\bar{p}, \vec{c}) = r_s I_s \max(0, \vec{r} \cdot \vec{c})^\alpha$$

where


$r_s$  - specular reflectance coefficient (how much light is reflected)

$I_s$  - “specular light source” (often =  $I_d$ )

$\vec{r}$  - direction of emission (normalized)



$\vec{c}$  - direction from the point to camera (normalized)

$\alpha$  - controls width of the highlight



# Specular Reflectance


- As we decrease alpha the reflection becomes more peaked (more like a mirror)




# Phong Reflectance Model

- Lets put all the peaces together
- **Remember:** light is additive

$$L(\bar{p}, \vec{c}) = r_d I_d \max(0, \vec{s} \cdot \vec{n}) + r_a I_a + r_s I_s \max(0, \vec{r} \cdot \vec{c})^\alpha$$



Diffuse



Specular




Ambient

# Phong Reflectance Model

- Lets put all the peaces together
- **Remember:** light is additive

$$L(\bar{p}, \vec{c}) = r_d I_d \max(0, \vec{s} \cdot \vec{n}) + r_a I_a + r_s I_s \max(0, \vec{r} \cdot \vec{c})^\alpha$$



# Phong Reflectance Model

- Lets put all the peaces together
- **Remember:** light is additive

$$L(\bar{\mathbf{p}}, \vec{\mathbf{c}}) = r_d \mathbf{I}_d \max(0, \vec{\mathbf{s}} \cdot \vec{\mathbf{n}}) + r_a \mathbf{I}_a + r_s \mathbf{I}_s \max(0, \vec{\mathbf{r}} \cdot \vec{\mathbf{c}})^\alpha$$

- for color

$$L_R(\bar{\mathbf{p}}, \vec{\mathbf{c}}) = r_{d,R} \mathbf{I}_{d,R} \max(0, \vec{\mathbf{s}} \cdot \vec{\mathbf{n}}) + r_{a,R} \mathbf{I}_{a,R} + r_{s,R} \mathbf{I}_{s,R} \max(0, \vec{\mathbf{r}} \cdot \vec{\mathbf{c}})^\alpha$$


$$L_G(\bar{\mathbf{p}}, \vec{\mathbf{c}}) = r_{d,G} \mathbf{I}_{d,G} \max(0, \vec{\mathbf{s}} \cdot \vec{\mathbf{n}}) + r_{a,G} \mathbf{I}_{a,G} + r_{s,G} \mathbf{I}_{s,G} \max(0, \vec{\mathbf{r}} \cdot \vec{\mathbf{c}})^\alpha$$

$$L_B(\bar{\mathbf{p}}, \vec{\mathbf{c}}) = r_{d,B} \mathbf{I}_{d,B} \max(0, \vec{\mathbf{s}} \cdot \vec{\mathbf{n}}) + r_{a,B} \mathbf{I}_{a,B} + r_{s,B} \mathbf{I}_{s,B} \max(0, \vec{\mathbf{r}} \cdot \vec{\mathbf{c}})^\alpha$$

- Notice we are still only considering point light source and are not considering secondary reflectance from surfaces

# Phong Reflectance Model


Increasing ratio of diffuse to  
specular reflection coefficients



Increasing exponent in specular term



# Ambient only



*Foley, van Dam, Feiner, Hughes, Plate II.28*

# Diffuse only



*Foley, van Dam, Feiner, Hughes, Plate II.29*




# Full Phong Model




*Foley, van Dam, Feiner, Hughes, Plate II.32*

# Multiple Light Sources



Blue Light Source



Blue+Red Light Sources



Blue+Red+Green  
Light Sources