

PROBABILISTIC GRAPHICAL MODELS
CPSC 532C (TOPICS IN AI)
STAT 521A (TOPICS IN MULTIVARIATE ANALYSIS)

LECTURE 2

Kevin Murphy

Wednesday 15 September, 2004

ADMINISTRIVIA

- Class web page http://www.cs.ubc.ca/~murphyk/Teaching/CS532c_Fall104/index.html
- Send email to 'majordomo@cs.ubc.ca' with the contents 'subscribe cpsc535c' to join class list.
(Note: email address does not correspond to correct class number!)
- Homework due in class on Monday 20th.
- Monday's class starts at 9.30am as usual.

REVIEW: PROBABILISTIC INFERENCE (STATE ESTIMATION)

- Inference is about estimating hidden (query) variables H from observed (visible) measurements v , which we can do as follows:


$$P(h|v) = \frac{P(v, h)}{\sum_{h'} P(v, h')}$$

- Examples:
 - Medical diagnosis: H diseases, v = findings/ symptoms,
 - Speech recognition: H = spoken words, v = acoustic waveform
 - Genetic pedigree analysis: H = genotype, v = phenotype

NAIVE INFERENCE

- Represent joint prob. distribution $P(C, S, R, W)$ as a 4D table of $2^4 = 32$ numbers.
- We observe the grass is wet and want to know how likely it was that the sprinkler caused this event.


$$\begin{aligned} P(s = 1 | w = 1) &= \frac{P(s = 1, w = 1)}{P(w = 1)} \\ &= \frac{\sum_{c=0}^1 \sum_{r=0}^1 P(s = 1, w = 1, R = r, C = c)}{\sum_{c,r,s} P(S = s, w = 1, R = r, C = c)} \end{aligned}$$



- Query/hidden vars = $\{S\}$, visible vars = $\{W\}$, nuisance vars = $\{C, R\}$.

NAIVE INFERENCE

- It is easy to marginalize a joint probability distribution when it is represented as a table
- e.g., $P(X, Y) = \sum_z P(X, Y, Z)$



GRAPHICAL MODELS

- Problems with representing joint as a big table
 - Representation: big table of numbers is hard to understand.
 - Inference: computing a marginal $P(X_i)$ takes $O(2^N)$ time.
 - Learning: there are $O(2^N)$ free parameters to estimate.
- Graphical models solve all 3 problems by providing a structured representation for joint probability distributions.
- Graphs encode conditional independence properties and represent families of probability distributions that satisfy these properties.
- Today we will study the relationship between graphs and independence properties.

INDEPENDENCE PROPERTIES OF DISTRIBUTIONS

- Defn: let $I(P)$ be the set of independence properties of the form $X \perp Y|Z$ that hold in distribution P .

X	Y	P(X,Y)
0	0	0.08
0	1	0.32
1	0	0.12
1	1	0.48

$$P(X = 1) = 0.48 + 0.12 = 0.6$$

$$P(Y = 1) = 0.32 + 0.48 = 0.8$$

$$P(X = 1, Y = 1) = 0.48 = 0.6 \times 0.8$$

$$P(X = x, Y = y) = P(X = x)P(Y = y) \forall x, y$$

$$\Rightarrow (X \perp Y) \in I(P)$$


$$\text{or } P \models (X \perp Y)$$

(LOCAL) INDEPENDENCE PROPERTIES OF DAGs



- Defn: let $I_l(G)$ be the set of local independence properties encoded by DAG G , namely:

$$\{X_i \perp \text{NonDescendants}(X_i) \mid \text{Parents}(X_i)\}$$



- i.e., a node is conditionally independent of its non-descendants given its parents.
- $\text{Ancestors}(X_i) \subseteq \text{NonDescendants}(X_i)$





EXAMPLE OF $I_l(G)$



G_0



$G_{X \rightarrow Y}$



$G_{Y \rightarrow X}$

$$I_l(G_\emptyset) = \{(X \perp Y)\}$$

$$I_l(G_{X \rightarrow Y}) = \emptyset$$

$$I_l(G_{Y \rightarrow X}) = \emptyset$$

I-MAPS

- Defn: A DAG G is an **I-map** (independence-map) of P if $I_l(G) \subseteq I(P)$.
- From previous example,

$$I_l(G_\emptyset) = \{(X \perp Y)\}$$

$$I_l(G_{X \rightarrow Y}) = \emptyset$$

$$I_l(G_{Y \rightarrow X}) = \emptyset$$

$$I(P) = \{(X \perp Y)\}$$

- Hence all three graphs are I-maps of P .

FROM I-MAP TO FACTORIZATION

- Defn: P factorizes according to G if P can be written as

$$P(X_1, \dots, X_N) = \prod_i P(X_i | \text{Pa}_G(X_i))$$

- Thm 3.2.6: If G is an I-map of P , then P factorizes according to G .
- Proof:


$$\begin{aligned} P(X_{1:N}) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots \text{ chain rule} \\ &= \prod_{i=1}^N P(X_i | X_{1:i-1}) \\ &= \prod_{i=1}^N P(X_i | \text{Pa}(X_i), \text{Ancestors}(X_i) \setminus \text{Pa}(X_i)) \\ &= \prod_{i=1}^N P(X_i | \text{Pa}(X_i)) \text{ since } G \text{ is I-map of } P \end{aligned}$$

BAYES NETS PROVIDE COMPACT REPRESENTATION OF JOINT PROBABILITY DISTRIBUTIONS

- Thm: If G is an I-map of P , then P factorizes according to G .
- Corollary: If G is an I-map of P , then we can represent P using G and a set of conditional probability distributions (CPDs), $P(X_i | \text{Pa}(X_i))$, one per node.
- Defn: A **Bayesian network** (aka **belief network**) representing distribution P is an I-map of P and a set of CPDs.
- For binary random variables, the Bayes net takes $O(N2^K)$ parameters ($K = \text{max. num. parents}$), whereas full joint takes $O(2^N)$ parameters.
- Factored representation is easier to understand, easier to learn and supports more efficient inference (see later lectures).

WATER SPRINKLER


$$P(X_{1:N}) = \prod_{i=1}^N P(X_i | \text{Pa}(X_i))$$



$$P(C, S, R, W) = P(C)P(S|C)P(R|C)P(W|S, R)$$

FROM FACTORIZATION TO I-MAP

- Thm 3.2.8: If P factorizes according to G , then G is an I-map of P .
- Proof: we must show $X \perp W|U$



$$\begin{aligned}
 P(X, W|U) &= \frac{P(X, W, U)}{P(U)} \\
 &= \frac{\sum_Y P(X, W, U, Y)}{P(U)} \\
 &= \frac{P(W)P(U|W)P(X|U) \sum_Y P(Y|X, W)}{P(U)} \\
 &= \frac{P(W, U)}{P(U)} P(X|U) \sum_Y P(Y|X, W) \\
 &= P(W|U)P(X|U)
 \end{aligned}$$

MINIMAL I-MAPS

- Let G be a fully connected DAG. Then $I_l(G) = \emptyset \subseteq I(P)$ for any P .
- Hence the complete graph is an I-map for any distribution.
- Defn: A DAG G is a **minimal I-map** for P if it is an I-map for P , and if the removal of even a single edge from G renders it not an I-map.
- Construction: pick a node ordering, then let the parents of node X_i be the minimal subset of $U \subseteq \{X_1, \dots, X_{i-1}\}$ s.t. $X_i \perp \{X_1, \dots, X_{i-1}\} \setminus U | U$.
- Defn (revised): A **Bayesian network** (aka **belief network**) representing distribution P is a *minimal* I-map of P and a set of CPDs.


GLOBAL MARKOV PROPERTIES OF DAGS

- By chaining together local independencies, we can infer more global independencies.
- Defn: X is **d-separated** (directed-separated) from Y given Z if along every undirected path between X and Y there is a node w s.t. either
 - W has converging arrows ($\rightarrow w \leftarrow$) and neither W nor its descendants are in Z ; or
 - W does not have converging arrows and $W \in Z$.
- Defn: $I(G) =$ all independence properties that correspond to d-separation:

$$I(G) = \{(X \perp Y | Z) : d\text{-sep}_G(X; Y | Z)\}$$

BAYES-BALL RULES

A is d-separated from B given C if we cannot send a ball from any node in A to any node in B according to the rules below, where shaded nodes are in C .



SOUNDNESS OF D-SEPARATION

- Thm 3.3.3 (**Soundness**): If P factorizes according to G , then $I(G) \subseteq I(P)$.
- i.e., any independence claim made by the graph is satisfied by all distributions P that factorize according to G (no false claims of independence).
- Pf: see later (when we discuss undirected graphs).

COMPLETENESS OF D-SEPARATION - v1

- Defn (Completeness) v1: For any distribution P that factorizes over G , if $(X \perp Y|Z) \in I(P)$, then $dsep_G(X; Y|Z)$.
- Contrapositive rule: $(A \Rightarrow B) \iff (\neg B \Rightarrow \neg A)$.
- Defn (Completeness, contrapositive form) v1. If X and Y are not d-separated given Z , then X and Y are dependent in all distributions P that factorize over G .
- This definition of completeness is too strong since P may have conditional independencies that are not evident from the graph.
- eg. Let G be the graph $X \rightarrow Y$, where $P(Y|X)$ is


A	$B = 0$	$B = 1$
0	0.4	0.6
1	0.4	0.6
- G is I-map of P since $I(G) = \emptyset \subseteq I(P) = \{(X \perp Y)\}$.
- But the CPD encodes $X \perp Y$ which is not evident in the graph.

COMPLETENESS OF D-SEPARATION - v2

- Defn (Completeness) v2: If $(X \perp Y|Z)$ in all distributions P that factorize over G , then $dsep_G(X; Y|Z)$.
- Defn (Completeness, contrapositive form) v2: If X and Y are not d-separated given Z , then X and Y are dependent in *some* distribution P that factorizes over G .
- Thm 3.3.5: d-separation is complete.
- Proof: See Koller & Friedman p90.
- Hence d-separation captures as many of the independencies as possible (without reference to the particular CPDs) for all distributions that factorize over some DAG.


P-MAPS

- Can we find a graph that captures all the independencies in an arbitrary distribution (and no more)?
- Defn: A DAG G is a **perfect map (P-map)** for a distribution P if $I(P) = I(G)$.
- Thm: not every distribution has a perfect map.
- Pf by counterexample. Suppose we have a model where $A \perp C | \{B, D\}$, and $B \perp D | \{A, C\}$. This cannot be represented by any Bayes net.
- e.g., BN1 wrongly says $B \perp D | A$, BN2 wrongly says $B \perp D$.



UNDIRECTED GRAPHICAL MODELS

- Graphs with one node per random variable and edges that connect pairs of nodes, but now the edges are undirected.
- Defn: Let H be an undirected graph. Then $sep_H(A; C|B)$ iff all paths between A and C go through some nodes in B (simple graph separation).




- Defn: the **global Markov properties** of a UG H are

$$I(H) = \{(X \perp Y|Z) : sep_H(X; Y|Z)\}$$

- UGs can model symmetric (non-causal) interactions that directed models cannot.
- aka Markov Random Fields, Markov Networks.

EXPRESSIVE POWER


- Can we always convert directed \leftrightarrow undirected?
- No.



(a)

No directed model
can represent these
and only these
independencies.

$$\mathbf{x} \perp \mathbf{y} \mid \{\mathbf{w}, \mathbf{z}\}$$
$$\mathbf{w} \perp \mathbf{z} \mid \{\mathbf{x}, \mathbf{y}\}$$



(b)

No undirected model
can represent these
and only these
independencies.

$$\mathbf{x} \perp \mathbf{y}$$

CONDITIONAL PARAMETERIZATION?

- In directed models, we started with $p(\mathbf{X}) = \prod_i p(\mathbf{x}_i | \mathbf{x}_{\pi_i})$ and we derived the d-separation semantics from that.
- Undirected models: have the semantics, need parametrization.
- What about this “conditional parameterization”?

$$p(\mathbf{X}) = \prod_i p(\mathbf{x}_i | \mathbf{x}_{\text{neighbours}(i)})$$

- Good: product of local functions.
Good: each one has a simple conditional interpretation.
Bad: local functions cannot be arbitrary, but must agree properly in order to define a valid distribution.

MARGINAL PARAMETERIZATION?

- OK, what about this “marginal parameterization”?

$$p(\mathbf{X}) = \prod_i p(\mathbf{x}_i, \mathbf{x}_{\text{neighbours}(i)})$$

- Good: product of local functions.
Good: each one has a simple marginal interpretation.
Bad: only very few pathological marginals on overlapping nodes can be multiplied to give a valid joint.



CLIQUE POTENTIALS

- Whatever factorization we pick, we know that only connected nodes can be arguments of a single local function.
- A *clique* is a fully connected subset of nodes.
- Thus, consider using a *product of clique potentials*:


$$P(\mathbf{X}) = \frac{1}{Z} \prod_{\text{cliques } c} \psi_c(\mathbf{x}_c) \qquad Z = \sum_{\mathbf{X}} \prod_{\text{cliques } c} \psi_c(\mathbf{x}_c)$$

- Each clique potential $\psi_c(\mathbf{x}_c) > 0$ is an arbitrary positive function of its arguments.
- The normalization term Z is called the partition function (a function of the parameters ψ) and ensures $\sum_{\mathbf{x}} P(\mathbf{x}) = 1$.
- Without loss of generality we can restrict ourselves to *maximal cliques*. (Why?)
- A distribution P that is representable by a UG H in this way is called a Gibbs distribution over H .

EXAMPLES OF CLIQUE POTENTIALS




(a)



(b)

INTERPRETATION OF CLIQUE POTENTIALS



- The model implies $\mathbf{x} \perp \mathbf{z} \mid \mathbf{y}$

$$p(\mathbf{x}, \mathbf{y}, \mathbf{z}) = p(\mathbf{y})p(\mathbf{x}|\mathbf{y})p(\mathbf{z}|\mathbf{y})$$

- We can write this as:

$$p(\mathbf{x}, \mathbf{y}, \mathbf{z}) = p(\mathbf{x}, \mathbf{y})p(\mathbf{z}|\mathbf{y}) = \psi_{\mathbf{xy}}(\mathbf{x}, \mathbf{y})\psi_{\mathbf{yz}}(\mathbf{y}, \mathbf{z})$$

$$p(\mathbf{x}, \mathbf{y}, \mathbf{z}) = p(\mathbf{x}|\mathbf{y})p(\mathbf{z}, \mathbf{y}) = \psi_{\mathbf{xy}}(\mathbf{x}, \mathbf{y})\psi_{\mathbf{yz}}(\mathbf{y}, \mathbf{z})$$

cannot have all potentials be marginals

cannot have all potentials be conditionals

- The positive clique potentials can only be thought of as general “compatibility”, “goodness” or “happiness” functions over their variables, but not as probability distributions.