

# CPSC 436D

## Video Game Programming



### *Transformations*




© Alla Sheffer

## COORDINATE SYSTEMS



*Coordinate system = Origin + Basis Vectors*




© Alla Sheffer



## COORDINATE SYSTEMS


*Coordinate system = Origin + Basis Vectors*



© Alla Sheffer



## COORDINATE SYSTEMS




$$P = O + x\vec{i} + y\vec{j}$$

equivalent:  $P = (x, y)$

© Alla Sheffer



## COORDINATE SYSTEMS



$$P = O + x\vec{i} + y\vec{j}$$

$\vec{F}_1$


$\vec{F}_2$

$\vec{F}_3$

© Alla Sheffer



## COORDINATE SYSTEMS



$$P = O + x\vec{i} + y\vec{j}$$

$\vec{F}_1$   $P(3,-1)$


$\vec{F}_2$

$\vec{F}_3$

© Alla Sheffer



## COORDINATE SYSTEMS



$$P = O + x\vec{i} + y\vec{j}$$

$$\mathbf{F}_1 \quad \mathbf{P}(3,-1)$$


$$\mathbf{F}_2 \quad \mathbf{P}(-1.5,2)$$

$$\mathbf{F}_3$$

© Alla Sheffer



## COORDINATE SYSTEMS



$$P = O + x\vec{i} + y\vec{j}$$

$$\mathbf{F}_1 \quad \mathbf{P}(3,-1)$$

$$\mathbf{F}_2 \quad \mathbf{P}(-1.5,2)$$

$$\mathbf{F}_3 \quad \mathbf{P}(1,2)$$


© Alla Sheffer



## Transformations

**Transformations as a change of frame**

$$P = O + xi\vec{i} + yj\vec{j}$$



check:  $P_1(3, -1)$  becomes  $P_2(-1.5, 2)$

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}_1 + x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix}_1 + y_1 \begin{bmatrix} 0 \\ 1 \end{bmatrix}_1$$

$$\begin{bmatrix} x \\ y \end{bmatrix}_2 = \begin{bmatrix} 0 \\ 3 \end{bmatrix}_2 + x_1 \begin{bmatrix} -0.5 \\ 0 \end{bmatrix}_2 + y_1 \begin{bmatrix} 0 \\ 1 \end{bmatrix}_2$$


$$\begin{bmatrix} x \\ y \end{bmatrix}_2 = \begin{bmatrix} -0.5 & 0 & 0 \\ 0 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_1$$

$$P_2 = MP_1$$

© Alla Sheffer

## TRANSFORMATIONS

**change of basis expressed using a matrix**



$$\begin{bmatrix} x \\ y \end{bmatrix}_2 = \begin{bmatrix} -0.5 & 0 & 0 \\ 0 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_1$$

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_2 = \begin{bmatrix} -0.5 & 0 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_1$$




© Alla Sheffer



## Usage of Transformations

set up the modeling matrix M

for each vertex v  
 $v' = Mv$



© Alla Sheffer




## Usage of Transformations

$$P = O + x\vec{i} + y\vec{j}$$

$$\begin{bmatrix} x_{obj} \\ y_{obj} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}_{obj} + x_{obj} \begin{bmatrix} 1 \\ 0 \end{bmatrix}_{obj} + y_{obj} \begin{bmatrix} 0 \\ 1 \end{bmatrix}_{obj}$$

$$\begin{bmatrix} x \\ y \end{bmatrix}_w = \begin{bmatrix} 3 \\ 1 \end{bmatrix}_w + x_{obj} \begin{bmatrix} 2 \\ 0 \end{bmatrix}_w + y_{obj} \begin{bmatrix} 0 \\ 2 \end{bmatrix}_w$$

$$\begin{bmatrix} x \\ y \end{bmatrix}_w = \begin{bmatrix} 2 & 0 & 3 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_{obj}$$



© Alla Sheffer



## Using Transformations

$$\begin{bmatrix} x \\ y \end{bmatrix}_2 = \begin{bmatrix} 2 & 0 & 3 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_1$$

2D


$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_w = \begin{bmatrix} 2 & 0 & 3 \\ 0 & 2 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}_{obj}$$

3D

$$\begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}_w = \begin{bmatrix} 2 & 0 & 0 & 3 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}_{obj}$$

© Alla Sheffer


## PIPELINE



© Alla Sheffer



## coordinate systems



© Alla Sheffer




## HOW TO TRANSFORM COORDINATES

- Between coordinate frames
- Or animate objects

© Alla Sheffer


## linear transformations



© Alla Sheffer

## Matrix representations

**Scale:**



© Alla Sheffer


## Matrix representations

**Scale:**

$$M = \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix}$$

Example:


$$\begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} \alpha \\ 2\beta \end{pmatrix}$$



© Alla Sheffer

## Matrix representations

**Rotation**



© Alla Sheffer


## Matrix representations

### Rotation

$$R(\alpha) = \begin{bmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{bmatrix}$$

Example:

$$\begin{bmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{bmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} \cos(\alpha) - \sin(\alpha) \\ \cos(\alpha) + \sin(\alpha) \end{pmatrix}$$



© Alla Sheffer

**What does this 2D transformation do?**

- A. Rotates by 90 deg
- B. Scales by a factor of 2
- C. Rotates by -90 deg
- D. Nothing

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

**What does this 2D transformation do?**

- A. Rotates by 90 deg
- B. Reflects the object
- C. Rotates by -90 deg
- D. Scales the object

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix},$$

© Alla Sheffer



## TRANSLATION

***There's a minor glitch.***

- Translation: can't be represented as 2x2 matrix multiplication

© Alla Sheffer



## general transformations

***We need to represent all the  
linear transformations + translation.***

***Ideas?***

$$T(\mathbf{v}) = M\mathbf{v} + \mathbf{b}$$

© Alla Sheffer



## AUGMENTED MATRIX

$$M_{2 \times 2} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x' \\ y' \end{bmatrix}$$

$$\begin{bmatrix} M_{2 \times 2} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix}$$

Haven't changed much, have we?

© Alla Sheffer



## AUGMENTED MATRIX

$$\begin{bmatrix} M_{2 \times 2} & b_x \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x' + b_x \\ y' + b_y \\ 1 \end{bmatrix} = \begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} + \mathbf{b}$$

Translation



© Alla Sheffer



## Affine transformations

- Linear (rotation, scaling, shear, reflections) + TRANSLATION

© Alla Sheffer



## Affine transformations

- Linear (rotation, scaling, shear, reflections) + TRANSLATION
- How to convert a linear transformation matrix into affine matrix?

© Alla Sheffer

## AFFINE Transformations


### Scale:

$$M = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$$

$$M = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example:

$$\begin{pmatrix} a \cdot 1 \\ b \cdot 2 \\ 1 \end{pmatrix} = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$$



© Alla Sheffer

## AFFINE Transformations


### Rotation

$$M = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$$

$$M = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example:

$$\begin{pmatrix} a \cdot 1 \\ b \cdot 2 \\ 1 \end{pmatrix} = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$$



© Alla Sheffer



## AFFINE Transformations

### Translation

$$M = \begin{bmatrix} 1 & 0 & C_x \\ 0 & 1 & C_y \\ 0 & 0 & 1 \end{bmatrix}$$

Example:

$$\begin{bmatrix} 1 & 0 & C_x \\ 0 & 1 & C_y \\ 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ 1 \end{pmatrix} = \begin{pmatrix} x + C_x \\ y + C_y \\ 1 \end{pmatrix}$$

© Alla Sheffer