



University of British Columbia  
CPSC 314 Computer Graphics  
Jan-Apr 2007

Tamara Munzner

**Math Review**

**Week 1, Wed Jan 10**




<http://www.ugrad.cs.ubc.ca/~cs314/Vjan2007>

# News

- signup sheet with name, email, program

# Review: Computer Graphics Defined



- CG uses
  - movies, games, art/design, ads, VR, visualization
- CG state of the art
  - photorealism achievable (in some cases)



# Correction: Expectations

- hard course!
  - heavy programming and heavy math
- fun course!
  - graphics programming addictive, create great demos
- programming prereq
  - CPSC 221 (Basic Algorithms and Data Structures)  
or CPSC 216 (Program Design and Data Structures)
  - course language is C++/C
- math prereq
  - MATH 200 (Calculus III)
  - MATH 221/223 (Matrix Algebra/Linear Algebra)

# Review: Rendering Capabilities



[www.siggraph.org/education/materials/HyperGraph/shutbug.htm](http://www.siggraph.org/education/materials/HyperGraph/shutbug.htm)

# Readings

- Mon (last time)
  - FCG Chap 1
- Wed (this time)
  - FCG Chap 2
    - except 2.5.1, 2.5.3, 2.7.1, 2.7.3, 2.8, 2.9, 2.11.
  - FCG Chap 5.1-5.2.5
    - except 5.2.3, 5.2.4
- Fri (next time)
  - RB Chap Introduction to OpenGL
  - RB Chap State Management and Drawing Geometric Objects
  - RB App Basics of GLUT (Aux in v 1.1)

# Today's Readings

- FCG Chapter 2: Miscellaneous Math
  - skim 2.2 (sets and maps), 2.3 (quadratic eqns)
  - important: 2.3 (trig), 2.4 (vectors), 2.5-6 (lines)  
2.10 (linear interpolation)
    - skip 2.5.1, 2.5.3, 2.7.1, 2.7.3, 2.8, 2.9
    - skip 2.11 now (covered later)
- FCG Chapter 5.1-5.25: Linear Algebra
  - skim 5.1 (determinants)
  - important: 5.2.1-5.2.2, 5.2.5 (matrices)
    - skip 5.2.3-4, 5.2.6-7 (matrix numerical analysis)

# Notation: Scalars, Vectors, Matrices

- scalar
  - (lower case, italic)
- vector
  - (lower case, bold)
- matrix
  - (upper case, bold)

$a$


$$\mathbf{a} = [a_1 \quad a_2 \quad \dots \quad a_n]$$

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$



# Vectors

- arrow: length and direction
  - oriented segment in nD space
- offset / displacement
  - location if given origin



# Column vs. Row Vectors

- row vectors  $\mathbf{a}_{row} = [a_1 \quad a_2 \quad \dots \quad a_n]$


- column vectors  $\mathbf{a}_{col} = \begin{bmatrix} a_1 \\ a_2 \\ \dots \\ a_n \end{bmatrix}$

- switch back and forth with transpose

$$\mathbf{a}_{col}^T = \mathbf{a}_{row}$$

# Vector-Vector Addition

- add: vector + vector = vector
- parallelogram rule
  - tail to head, complete the triangle



algebraic

$$\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_1 + v_1 \\ u_2 + v_2 \\ u_3 + v_3 \end{bmatrix}$$


examples:

$$(3,2) + (6,4) = (9,6)$$
$$(2,5,1) + (3,1,-1) = (5,6,0)$$

# Vector-Vector Subtraction

- subtract: vector - vector = vector

$$\mathbf{u} - \mathbf{v} = \begin{bmatrix} u_1 - v_1 \\ u_2 - v_2 \\ u_3 - v_3 \end{bmatrix}$$




$$(3,2) - (6,4) = (-3,-2)$$

$$(2,5,1) - (3,1,-1) = (-1,4,2)$$

# Vector-Vector Subtraction

- subtract: vector - vector = vector



$$\mathbf{u} - \mathbf{v} = \begin{bmatrix} u_1 - v_1 \\ u_2 - v_2 \\ u_3 - v_3 \end{bmatrix}$$



$$(3,2) - (6,4) = (-3,-2)$$


$$(2,5,1) - (3,1,-1) = (-1,4,2)$$

argument reversal



# Scalar-Vector Multiplication

- multiply: scalar \* vector = vector
  - vector is scaled



$$a * \mathbf{u} = (a * u_1, a * u_2, a * u_3)$$

$$2 * (3, 2) = (6, 4)$$

$$.5 * (2, 5, 1) = (1, 2.5, .5)$$

# Vector-Vector Multiplication

- multiply: vector \* vector = scalar
- dot product, aka inner product

$$\mathbf{u} \bullet \mathbf{v}$$

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} \bullet \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = (u_1 * v_1) + (u_2 * v_2) + (u_3 * v_3)$$

# Vector-Vector Multiplication

- multiply: vector \* vector = scalar
- dot product, aka inner product

$$\mathbf{u} \bullet \mathbf{v}$$

$$\begin{array}{|c|c|} \hline u_1 & v_1 \\ \hline u_2 & v_2 \\ \hline u_3 & v_3 \\ \hline \end{array} = (u_1 * v_1) + (u_2 * v_2) + (u_3 * v_3)$$



# Vector-Vector Multiplication


- multiply: vector \* vector = scalar
- dot product, aka inner product

$$\mathbf{u} \bullet \mathbf{v}$$

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} \bullet \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = (u_1 * v_1) + (u_2 * v_2) + (u_3 * v_3)$$

- geometric interpretation
  - lengths, angles
  - can find angle between two vectors

$$\mathbf{u} \bullet \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta$$




# Dot Product Geometry


- can find length of projection of  $\mathbf{u}$  onto  $\mathbf{v}$

$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta$$

$$\|\mathbf{u}\| \cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|}$$



- as lines become perpendicular,  $\mathbf{u} \cdot \mathbf{v} \rightarrow 0$



## Dot Product Example

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} \cdot \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = (u_1 * v_1) + (u_2 * v_2) + (u_3 * v_3)$$

$$\begin{bmatrix} 6 \\ 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 7 \\ 3 \end{bmatrix} = (6 * 1) + (1 * 7) + (2 * 3) = 6 + 7 + 6 = 19$$

# Vector-Vector Multiplication, The Sequel

- multiply: vector \* vector = vector
- cross product
  - algebraic

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} \times \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} u_2 v_3 - u_3 v_2 \\ u_3 v_1 - u_1 v_3 \\ u_1 v_2 - u_2 v_1 \end{bmatrix}$$

# Vector-Vector Multiplication, The Sequel

- multiply: vector \* vector = vector
- cross product
  - algebraic

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} \times \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} u_2 v_3 - u_3 v_2 \\ u_3 v_1 - u_1 v_3 \\ u_1 v_2 - u_2 v_1 \end{bmatrix}$$

# Vector-Vector Multiplication, The Sequel

- multiply: vector \* vector = vector
- cross product
  - algebraic

$$\begin{array}{l} 3 \\ 1 \\ 2 \end{array} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} \times \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} u_2 v_3 - u_3 v_2 \\ u_3 v_1 - u_1 v_3 \\ u_1 v_2 - u_2 v_1 \end{bmatrix}$$


# Vector-Vector Multiplication, The Sequel

- multiply: vector \* vector = vector
- cross product
  - algebraic
  - geometric

$$\|\mathbf{a} \times \mathbf{b}\| = \|\mathbf{a}\| \|\mathbf{b}\| \sin \theta$$


- $\|\mathbf{a} \times \mathbf{b}\|$  parallelogram area
- $\mathbf{a} \times \mathbf{b}$  perpendicular to parallelogram

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} \times \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} u_2 v_3 - u_3 v_2 \\ u_3 v_1 - u_1 v_3 \\ u_1 v_2 - u_2 v_1 \end{bmatrix}$$



# RHS vs. LHS Coordinate Systems


- right-handed coordinate system **convention**



right hand rule:  
index finger x, second finger y;  
right thumb points up

$$\mathbf{Z} = \mathbf{X} \times \mathbf{y}$$

- left-handed coordinate system



left hand rule:  
index finger x, second finger y;  
left thumb points down


$$\mathbf{Z} = \mathbf{X} \times \mathbf{y}$$



# Basis Vectors

- take any two vectors that are **linearly independent** (nonzero and nonparallel)
  - can use linear combination of these to define any other vector:

$$\mathbf{c} = w_1 \mathbf{a} + w_2 \mathbf{b}$$




# Orthonormal Basis Vectors

- if basis vectors are **orthonormal** (**orthogonal** (mutually perpendicular) and unit length)
  - we have Cartesian coordinate system
  - familiar Pythagorean definition of distance

orthonormal algebraic properties


$$\|\mathbf{x}\| = \|\mathbf{y}\| = 1,$$

$$\mathbf{x} \cdot \mathbf{y} = 0$$




# Basis Vectors and Origins

- **coordinate system**: just basis vectors
  - can only specify offset: vectors
- **coordinate frame**: basis vectors and origin
  - can specify location as well as offset: points



$$\mathbf{p} = \mathbf{o} + x\mathbf{i} + y\mathbf{j}$$


# Working with Frames



$$\mathbf{p} = \mathbf{o} + x\mathbf{i} + y\mathbf{j}$$

$F_1$


# Working with Frames



$$\mathbf{p} = \mathbf{o} + x\mathbf{i} + y\mathbf{j}$$

$$F_1 \quad \mathbf{p} = (3, -1)$$

# Working with Frames




$$\mathbf{p} = \mathbf{0} + x\mathbf{i} + y\mathbf{j}$$

$$F_1 \quad \mathbf{p} = (3, -1)$$

$$F_2$$

# Working with Frames




$$\mathbf{p} = \mathbf{o} + x\mathbf{i} + y\mathbf{j}$$

$$F_1 \quad \mathbf{p} = (3, -1)$$

$$F_2 \quad \mathbf{p} = (-1.5, 2)$$

# Working with Frames



$$\mathbf{p} = \mathbf{o} + x\mathbf{i} + y\mathbf{j}$$


$$F_1 \quad \mathbf{p} = (3, -1)$$

$$F_2 \quad \mathbf{p} = (-1.5, 2)$$

$$F_3$$



# Working with Frames



$$\mathbf{p} = \mathbf{o} + x\mathbf{i} + y\mathbf{j}$$

$$F_1 \quad \mathbf{p} = (3, -1)$$

$$F_2 \quad \mathbf{p} = (-1.5, 2)$$


$$F_3 \quad \mathbf{p} = (1, 2)$$

# Named Coordinate Frames

- origin and basis vectors  $\mathbf{p} = \mathbf{o} + a\mathbf{x} + b\mathbf{y} + c\mathbf{z}$
- pick canonical frame of reference
  - then don't have to store origin, basis vectors
  - just  $\mathbf{p} = (a, b, c)$
  - convention: Cartesian orthonormal one on previous slide
- handy to specify others as needed
  - airplane nose, looking over your shoulder, ...
  - really common ones given names in CG
    - object, world, camera, screen, ...


# Lines

- slope-intercept form
  - $y = mx + b$
- implicit form
  - $y - mx - b = 0$
  - $Ax + By + C = 0$
  - $f(x,y) = 0$



# Implicit Functions

- find where function is 0
  - plug in  $(x,y)$ , check if
    - 0: on line
    - $< 0$ : inside
    - $> 0$ : outside
- analogy: terrain
  - sea level:  $f=0$
  - altitude: function value
  - topo map: equal-value contours (level sets)



# Implicit Circles

- $f(x, y) = (x - x_c)^2 + (y - y_c)^2 - r^2$ 
  - circle is points  $(x, y)$  where  $f(x, y) = 0$
- $p = (x, y), c = (x_c, y_c) : (\mathbf{p} - \mathbf{c}) \cdot (\mathbf{p} - \mathbf{c}) - r^2 = 0$ 
  - points  $\mathbf{p}$  on circle have property that vector from  $\mathbf{c}$  to  $\mathbf{p}$  dotted with itself has value  $r^2$
- $\|\mathbf{p} - \mathbf{c}\|^2 - r^2 = 0$ 
  - points  $\mathbf{p}$  on the circle have property that squared distance from  $\mathbf{c}$  to  $\mathbf{p}$  is  $r^2$
- $\|\mathbf{p} - \mathbf{c}\| - r = 0$ 
  - points  $\mathbf{p}$  on circle are those a distance  $r$  from center point  $\mathbf{c}$


# Parametric Curves

- parameter: index that changes continuously
  - $(x,y)$ : point on curve
  - $t$ : parameter
- vector form
  - $\mathbf{p} = f(t)$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} g(t) \\ h(t) \end{bmatrix}$$

## 2D Parametric Lines

- $$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x_0 + t(x_1 - x_0) \\ y_0 + t(y_1 - y_0) \end{bmatrix}$$
- $\mathbf{p}(t) = \mathbf{p}_0 + t(\mathbf{p}_1 - \mathbf{p}_0)$
- $\mathbf{p}(t) = \mathbf{o} + t(\mathbf{d})$
- start at point  $\mathbf{p}_0$ ,  
go towards  $\mathbf{p}_1$ ,  
according to parameter  $t$ 
  - $\mathbf{p}(0) = \mathbf{p}_0$ ,  $\mathbf{p}(1) = \mathbf{p}_1$



# Linear Interpolation

- parametric line is example of general concept
  - $\mathbf{p}(t) = \mathbf{p}_0 + t(\mathbf{p}_1 - \mathbf{p}_0)$
  - interpolation
    - $\mathbf{p}$  goes through  $\mathbf{a}$  at  $t = 0$
    - $\mathbf{p}$  goes through  $\mathbf{b}$  at  $t = 1$
  - linear
    - weights  $t, (1-t)$  are linear polynomials in  $t$



# Matrix-Matrix Addition

- add: matrix + matrix = matrix

$$\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} + \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} = \begin{bmatrix} n_{11} + m_{11} & n_{12} + m_{12} \\ n_{21} + m_{21} & n_{22} + m_{22} \end{bmatrix}$$

- example

$$\begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} -2 & 5 \\ 7 & 1 \end{bmatrix} = \begin{bmatrix} 1 + (-2) & 3 + 5 \\ 2 + 7 & 4 + 1 \end{bmatrix} = \begin{bmatrix} -1 & 8 \\ 9 & 5 \end{bmatrix}$$

# Scalar-Matrix Multiplication

- multiply: scalar \* matrix = matrix

$$a \begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} = \begin{bmatrix} a * m_{11} & a * m_{12} \\ a * m_{21} & a * m_{22} \end{bmatrix}$$

- example

$$3 \begin{bmatrix} 2 & 4 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 3 * 2 & 3 * 4 \\ 3 * 1 & 3 * 5 \end{bmatrix} = \begin{bmatrix} 6 & 12 \\ 3 & 15 \end{bmatrix}$$

# Matrix-Matrix Multiplication

- can only multiply (n,k) by (k,m):  
number of left cols = number of right rows

- legal

$$\begin{bmatrix} a & b & c \\ e & f & g \end{bmatrix} \begin{bmatrix} h & i \\ j & k \\ l & m \end{bmatrix}$$

- undefined

$$\begin{bmatrix} a & b & c \\ e & f & g \\ o & p & q \end{bmatrix} \begin{bmatrix} h & i \\ j & k \end{bmatrix}$$

# Matrix-Matrix Multiplication

- row by column

$$\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} = \begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}$$

$$p_{11} = m_{11}n_{11} + m_{12}n_{21}$$

# Matrix-Matrix Multiplication

- row by column

$$\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} = \begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}$$

$$p_{11} = m_{11}n_{11} + m_{12}n_{21}$$

$$p_{21} = m_{21}n_{11} + m_{22}n_{21}$$

# Matrix-Matrix Multiplication

- row by column

$$\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} = \begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}$$

$$p_{11} = m_{11}n_{11} + m_{12}n_{21}$$

$$p_{21} = m_{21}n_{11} + m_{22}n_{21}$$

$$p_{12} = m_{11}n_{12} + m_{12}n_{22}$$

# Matrix-Matrix Multiplication

- row by column

$$\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} = \begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}$$

$$p_{11} = m_{11}n_{11} + m_{12}n_{21}$$

$$p_{21} = m_{21}n_{11} + m_{22}n_{21}$$

$$p_{12} = m_{11}n_{12} + m_{12}n_{22}$$

$$p_{22} = m_{21}n_{12} + m_{22}n_{22}$$

# Matrix-Matrix Multiplication

- row by column

$$\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} = \begin{bmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{bmatrix}$$

$$p_{11} = m_{11}n_{11} + m_{12}n_{21}$$

$$p_{21} = m_{21}n_{11} + m_{22}n_{21}$$

$$p_{12} = m_{11}n_{12} + m_{12}n_{22}$$

$$p_{22} = m_{21}n_{12} + m_{22}n_{22}$$

- noncommutative: **AB**  $\neq$  **BA**



# Matrix-Vector Multiplication

- points as column vectors: postmultiply

$$\begin{bmatrix} x' \\ y' \\ z' \\ h' \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \\ m_{41} & m_{42} & m_{43} & m_{44} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ h \end{bmatrix} \quad \mathbf{p}' = \mathbf{M}\mathbf{p}$$

- points as row vectors: premultiply

$$\begin{bmatrix} x' & y' & z' & h' \end{bmatrix} = \begin{bmatrix} x & y & z & h \end{bmatrix} \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \\ m_{41} & m_{42} & m_{43} & m_{44} \end{bmatrix}^T \quad \mathbf{p}'^T = \mathbf{p}^T \mathbf{M}^T$$

# Matrices

- transpose  $\begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \\ m_{41} & m_{42} & m_{43} & m_{44} \end{bmatrix}^T = \begin{bmatrix} m_{11} & m_{21} & m_{31} & m_{41} \\ m_{12} & m_{22} & m_{32} & m_{42} \\ m_{13} & m_{23} & m_{33} & m_{43} \\ m_{14} & m_{24} & m_{34} & m_{44} \end{bmatrix}$

- identity  $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

- inverse  $\mathbf{AA}^{-1} = \mathbf{I}$

- not all matrices are invertible

# Matrices and Linear Systems

- linear system of n equations, n unknowns

$$3x + 7y + 2z = 4$$

$$2x - 4y - 3z = -1$$

$$5x + 2y + z = 1$$

- matrix form  **$Ax=b$**

$$\begin{bmatrix} 3 & 7 & 2 \\ 2 & -4 & -3 \\ 5 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ -1 \\ 1 \end{bmatrix}$$